Colonoscopy Prep (ages 1-3):

WHAT YOU NEED:

• One large bottle of **polyethylene glycol 3350**. The most common brand is MiraLAX® but ClearLAX, Powderlax, etc. are examples of generic store brands.





EXAMPLES OF POLYETHLENE GLYCOL 3350 PRODUCTS

HOW TO USE MiraLAX®:

- MiraLAX is a stool softener for children ages 6 months and older. It works by bringing more water into the colon and softening the stool, making it easier to pass.
- MiraLAX comes in a powder that is mixed in clear liquid.
- One dose = 1 capful of powder + 8 ounces (1 cup) of one of the following clear liquids:
 - Water
 - Flavored water
 - Jell-O® gelatin
 - Slushies
 - Icee®

- Popsicles
- Gatorade®
- Powerade®
- Pedialyte®
- Kool-Aid®

- Sprite[®]
- Starry
- Apple juice
- White grape juice
- Chicken bouillon cubes (not canned)







AVOID:

- All solid foods
- Milk or dairy products
- Red- or purple-colored liquids
- Orange juice or juice with pulp
- Gatorade "Recovery" brand, which contains whey protein that should be avoided during prep

The goal of the bowel prep is clear stool from your child's intestine so we can safely see and navigate during the procedure.

- During the prep your child's stool should become light yellow to clear in color.
- Your child needs to stay hydrated while doing their prep.
 Encourage extra clear liquids!
- Poor bowel prep will result in rescheduling the procedure.

LEARN MORE



PLEASE FOLLOW THIS SCHEDULE

At 8 a.m. (one day before procedure)

- Your child may eat a light breakfast (for example, pancakes, French toast, biscuit, cereal, etc.)
- Begin clear liquid diet (see list of acceptable clear liquids). No solid food should be eaten.
- Your child can have as much clear liquid as they want – this is important to help keep your child hydrated.
- Give one dose of MiraLAX

Next: Give one dose of MiraLAX each hour as scheduled below

□ 9 a.m.	☐ 2 p.m. Please call Gl Procedure at 816-234-3704 if your child has not yet passed stool.
□ 10 a.m.	
□ 11 a.m.	
□ 12 p.m.	
□ 1 p.m.	□ 3 p.m.
	□ 4 p.m.

If your child's stools are clear or light yellow, they are ready for their procedure.

□ 5 p.m.

If stools are not clear or light yellow continue giving more hourly doses of MiraLAX.

ADDITIONAL INFORMATION

If you have questions, call 816-234-3000 and ask for the GI doctor on call for recommendations.



Copyright © 2023 The Children's Mercy Hospital. All rights reserved. 24-GI-2000 12/23