

Behavioral Escalation in the ED Clinical Pathway Synopsis


Behavioral Escalation in the ED (Initial Presentation)

Inclusion Criteria:
• Agitation - Restlessness, pacing, hand ringing, etc.
• Aggression - Change in body language, argumentative or defensive, verbal or physical threats
• Self Harm Behavior - Expressed thought to harm self/others, present injurious behaviors or plan
• Threat to Harm Others - Expressed thought to harm others, present threatening behaviors
• Elopement - Verbal request to leave, movement towards exit, physical attempt to leave
Exclusion Criteria:
• Any non-CM patient

Sources of Escalating Behavior
• Pain
• Hunger
• Environmental factors
• Delirium
• Substance abuse or withdrawal <ul style="list-style-type: none"> ◦ Consider nicotine withdrawal (refer to Nicotine Cessation Clinical Pathway) ◦ Consider opioid withdrawal (refer to Opioid Withdrawal Treatment: Buprenorphine/Naloxone, (Suboxone) Clinical Pathway) ◦ Consider Toxicology Consult
• Developmental delay or autism
• Psychiatric diagnoses
(Source of Behavioral Escalation Management and Medications)

Indications for Restraints
• First line of care should be seclusion (if available) before medications or restraints.
• If seclusion fails (risk of self-harm or while in seclusion) proceed to medications and/or restraint as needed (See Restraint or Seclusion Policy).

Associated CM Policies:
• Behavior Management
• Internal Transport of Disruptive and Flight Risk Patients
• Patient Transport within the Facility Guidelines
• Restraint or Seclusion
• Continuous Patient Observation


QR code
for mobile
view

These clinical pathways do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare a clinical pathway for each. Accordingly, these clinical pathways should guide care with the understanding that departures from them may be required at times.

**Behavioral Escalation in the ED
(Huddle with Care Team)**

Sources of Escalating Behavior

- Pain
- Hunger
- Environmental factors
- Delirium
- Substance abuse or withdrawal
 - Consider nicotine withdrawal (refer to [Nicotine Cessation Clinical Pathway](#))
 - Consider opioid withdrawal (refer to [Opioid Withdrawal Treatment: Buprenorphine/Naloxone \[Suboxone\] Clinical Pathway](#))
 - Consider Toxicology Consult
- Developmental delay or autism
- Psychiatric diagnoses

[\(Source of Behavioral Escalation Management and Medications\)](#)



Huddle with Care Team

- Involve family when possible
- Verify and administer home medications
- Have security at bedside, if needed
- Collaborate with Social Work, House Shift Supervisor, Child Life and Staff Advocacy Team, as needed
- Discuss whether or not 1:1 is needed

Quick Access:
[Source of Behavioral Escalation Management and Medications](#)
[POPS \(Parents Offering Parent Support\)](#)


Indications for Restraints

- First line of care should be seclusion (if available) **before** medications or restraints.
- If seclusion fails (risk of self-harm or while in seclusion) proceed to medications and/or restraint as needed (See [Restraint or Seclusion Policy](#)).

Consider the following as needed, if not already done:

- 1:1 observation (See, [Continuous Patient Observation Policy](#))
- Move to Safer Room, if possible
- Seclusion
 - Consider physical restraints if seclusion has been unsuccessful (See [Restraint or Seclusion Policy](#))
- Administration of [behavioral medications based on source](#)

Note: If prior visits, review notes for previous behavioral plan, effective strategies, or medication plans




QR code
for mobile view

Associated CMKC Policies:

- Behavior Management
- Internal Transport of Disruptive and Flight Risk Patients
- Patient Transport within the Facility Guidelines
- Restraint or Seclusion
- Continuous Patient Observation

These clinical pathways do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare a clinical pathway for each. Accordingly, these clinical pathways should guide care with the understanding that departures from them may be required at times.



These clinical pathways do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare a clinical pathway for each. Accordingly, these clinical pathways should guide care with the understanding that departures from them may be required at times.

Table of Contents

Objective of Clinical Pathway.....	5
Background	5
Target Users.....	5
Target Population.....	5
Practice Recommendations	5
Additional Questions Posed by the Clinical Pathway Committee.....	5
Updates from Previous Versions of the Clinical Pathway	5
Recommendation Specific for Children's Mercy	6
Measures	6
Value Implication	6
Organizational Barriers and Facilitators	6
Diversity/Equity/Inclusion	6
Clinical Pathway Preparation.....	6
Clinical Pathway Committee Members and Representation.....	6
Clinical Pathway Development Funding	7
Approval Process.....	7
Review Requested	7
Version History	7
Date for Next Review.....	7
Implementation & Follow-Up	7
Disclaimer	8
References	9

These clinical pathways do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare a clinical pathway for each. Accordingly, these clinical pathways should guide care with the understanding that departures from them may be required at times.

Objective of Clinical Pathway

The objective of the Behavioral Escalation in the ED Clinical Pathway is to provide a structured, evidence-based approach to managing children and adolescents who exhibit challenging or aggressive behaviors. These pathways are designed to ensure that healthcare providers follow consistent, safe, and effective de-escalation, diagnosis, and treatment strategies while promoting a compassionate, patient-centered environment.

Background

Pediatric behavioral escalation presents significant challenges in the emergency department (ED) (Saidinejad et al., 2023). Behavioral crises in children, marked by aggressive or disruptive actions, are becoming more common as mental health-related visits to the ED increase (Radhakrishnan, 2022). A substantial proportion of healthcare staff, particularly emergency nurses, have experienced incidents of verbal or physical abuse in these situations (Benning et al., 2024). In pediatric cases, mental health visits frequently escalate to a point of requiring interventions such as physical restraints, with 6% to 10% of these cases involving such measures (Gerson et al., 2019). Implementing effective management strategies, including prevention, early identification of escalation, verbal de-escalation techniques, and timely behavioral interventions, are critical to maintaining a safe environment for patients and staff (Gerson et al., 2019).

Target Users

- Physicians (Emergency Medicine, Fellows, Resident Physicians)
- Nurse Practitioners
- Nurses
- Social Work
- Care Assistance

Target Population

Inclusion Criteria

- Patients presenting or developing the following behaviors
 - **Aggression** - Change in body language, argumentative or defensive, verbal or physical escalation
 - **Elopement** - Verbal request to leave, movement towards exit, physical attempt to leave
 - **Self-Harm Behavior** - Expressed thought to harm self/others, present injurious behaviors or plan, intent during care
 - **Threat to Harm Others** - Expressed thought to harm others, present plan or intent during care

Exclusion Criteria

- Any non-Children's Mercy Kansas City patients
 - Including parents and family members

Practice Recommendations

Please refer to the American Association for Emergency Psychiatry's Best Practices for Evaluation and Treatment of Agitated Children and Adolescents (BETA) for full practice recommendations, evaluation, and treatment recommendations (Gerson et al., 2019).

Additional Questions Posed by the Clinical Pathway Committee

No additional clinical questions beyond the scope of Emergency Psychiatry's consensus statement were posed for this review.

Updates from Previous Versions of the Clinical Pathway

- Renamed pathway from "Aggressive Patient in ED" to "Behavioral Escalation in the ED"
- Broadened scope to include not only aggression, but also concern for self-harm or elopement
- Removed the Behavioral Health Observation Tool and instead incorporated key points into the pathway itself
- Recategorized level of aggression from mild/moderate/severe to immediate risk versus potential risk
- Added recommendations for keeping the patient care environment safe for patients and medical team members

These clinical pathways do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare a clinical pathway for each. Accordingly, these clinical pathways should guide care with the understanding that departures from them may be required at times.

- Highlighted behavioral strategies for de-escalation
- Added the use of seclusion as a recommended intervention
- Recommend pharmacologic treatment based on source of behavioral escalation
- Expanded medication table, including dose, route, time of onset, and time of peak effect

Recommendation Specific for Children's Mercy

Children's Mercy adopted the majority of the practice recommendations made by American Association for Emergency Psychiatry's Best Practices for Evaluation and Treatment of Agitated Children and Adolescents (BETA) for full practice recommendations, evaluation, and treatment recommendations (Gerson et al., 2019).

Variations/Additions include: No deviations were made from the practice recommendation paper, but logistical processes specific to Children's Mercy were added.

Measures

- Use of the Behavioral Escalation Clinical Pathway

Value Implication

- Improved safety of patients
- Improved safety of medical team members
- Decreased need for physical restraints
- Decreased unwarranted variation in care

Organizational Barriers and Facilitators**Potential Barriers**

- Variability in training in de-escalation strategies by medical team members
- Variability in resources based on location (availability of Safe Rooms)

Potential Facilitators

- Collaborative engagement across care continuum settings during clinical pathway development
- High rate of use of clinical pathways

Diversity/Equity/Inclusion

Our aim is to provide equitable care. These issues were discussed with the Committee, reviewed in the literature, and discussed prior to making any practice recommendations.

Associated Policies

- Behavior Management
- Internal Transport of Disruptive and Flight Risk Patients
- Patient Transport within the Facility Guidelines
- Restraint or Seclusion
- Continuous Patient Observation

Clinical Pathway Preparation

This pathway was prepared by the Evidence Based Practice (EBP) Department in collaboration with the Behavioral Escalation Clinical Pathway Committee composed of content experts at Children's Mercy Kansas City. If a conflict of interest is identified, the conflict will be disclosed next to the committee member's name.

Clinical Pathway Committee Members and Representation

- John Graham, MD | Emergency Department | Committee Chair
- Dani Wentzel, DO | Developmental and Behavioral Health | Committee Chair
- Christopher Kennedy, MD | Emergency Department | Committee Member
- Brandi Missel, ARNP | Emergency Department | Committee Member
- Alec Bernstein, PhD, BCBA-D, LBA | Developmental and Behavioral Health | Committee Member
- Ibad Siddiqi, PharmD | Pharmacy | Committee Member

These clinical pathways do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare a clinical pathway for each. Accordingly, these clinical pathways should guide care with the understanding that departures from them may be required at times.

- Emily Snow, LMSW, MHSA | Developmental and Behavioral Health | Committee Member
- Philip Lawler, MBA, CPP | Security/Transportation | Committee Member
- Katie Stangler, MSN, APRN, CPNP, CCRN | Emergency and Trauma Services | Committee Member
- Tanis Stewart, MSN, RN, FNP-BC, CPN | Kansas Emergency Department | Committee Member
- Erin Bass, LCSW, LSCSW | Social Work | Committee Member
- Kathryn Worland, MSW, LCSW, LSCSW | Social Work | Committee Member
- Kourtney Frederes, RN | Emergency Department | Committee Member
- Holly Reid, RN | Emergency Department | Committee Member
- Michele Smith, CNA | Emergency Department | Committee Member
- Sarah Nienhaus, MSN, RN, CPEN | Emergency Department | Committee Member
- Viktoriya Stoycheva, MHA, BSN, RN, CPN | Emergency Department | Committee Member

EBP Committee Members

- Kathleen Berg, MD, FAAP | Hospitalist, Evidence Based Practice
- Jarrod Dusin, MS, RD, LD, CPHQ | Evidence Based Practice

Clinical Pathway Development Funding

The development of this clinical pathway was underwritten by the following departments/divisions: Evidence Based Practice, Emergency Department, Developmental and Behavioral Health, and Social Work.

Conflict of Interest

The contributors to the Behavioral Escalation Clinical Pathway have no conflicts of interest to disclose related to the subject matter or materials discussed.

Approval Process

- This pathway was reviewed and approved by the Behavioral Escalation in the ED Clinical Pathway Committee, Content Expert Departments/Divisions, and the EBP Department; after which they were approved by the Medical Executive Committee.
- Pathways are reviewed and updated as necessary every 3 years within the EBP Department at CMKC. Content expert teams are involved with every review and update.

Review Requested

Department/Unit	Date Obtained
Emergency Department	February 2025
Developmental and Behavioral Health	October 2024
Security and Transportation	October 2024
Evidence Based Practice	October 2024
Pharmacy	October 2024
Social Work	October 2024

Version History

Date	Comments
7/23/2021	Version one – The pathway was developed to manage aggressive patients in the ED using a behavioral health observation tool (BHOT).
3/2025	Version two – Renamed from aggressive patients in the ED, BHOT removed, scope of patients expanded, treatment based on source of behavior, medication table expanded.

Date for Next Review

- March 2028

Implementation & Follow-Up

- Education was provided to all stakeholders:

These clinical pathways do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare a clinical pathway for each. Accordingly, these clinical pathways should guide care with the understanding that departures from them may be required at times.

Nursing units where the clinical pathway is used
Providers from the Emergency Department and Developmental and Behavioral Health

- Additional institution-wide announcements were made via email, hospital website, and relevant huddles.

Disclaimer

When evidence is lacking or inconclusive, options in care are provided in the supporting documents and the power plan(s) that accompany the clinical pathway.

These clinical pathways do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time.

It is impossible to anticipate all possible situations that may exist and to prepare clinical pathways for each. Accordingly, these clinical pathways should guide care with the understanding that departures from them may be required at times.

These clinical pathways do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare a clinical pathway for each. Accordingly, these clinical pathways should guide care with the understanding that departures from them may be required at times.

References

Benning, L., Teepe, G. W., Kleinekort, J., Thoma, J., Röttger, M. C., Prunotto, A., ... & Hans, F. P. (2024). Workplace violence against healthcare workers in the emergency department—a 10-year retrospective single-center cohort study. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine*, 32(1), 88.

Gerson, R., Malas, N., Feuer, V., Silver, G. H., Prasad, R., & Mroczkowski, M. M. (2019). Best practices for evaluation and treatment of agitated children and adolescents (BETA) in the emergency department: consensus statement of the American Association for Emergency Psychiatry. *Western Journal of Emergency Medicine*, 20(2), 409.

Radhakrishnan, L. (2022). Pediatric emergency department visits associated with mental health conditions before and during the COVID-19 pandemic—United States, January 2019–January 2022. *MMWR. Morbidity and mortality weekly report*, 71.

Saidinejad, M., Duffy, S., Wallin, D., Hoffmann, J. A., Joseph, M. M., Schieferle Uhlenbrock, J., ... & EMERGENCY NURSES ASSOCIATION Pediatric Committee. (2023). The management of children and youth with pediatric mental and behavioral health emergencies. *Pediatrics*, 152(3).

These clinical pathways do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare a clinical pathway for each. Accordingly, these clinical pathways should guide care with the understanding that departures from them may be required at times.