

**Dosing of antibiotics found in algorithm:**

- Amoxicillin 40-50 mg/kg/dose PO BID (max 2000 mg/dose)
- Amoxicillin/clavulanate 40-50 mg/kg/dose (amoxicillin component) PO BID (max 2000 mg amoxicillin component /dose)
- Cefuroxime 250 mg PO BID (max 500 mg/dose) for children able to swallow pills (only available in tablet form)
- Cefdinir 7 mg/kg/dose PO BID (max 300 mg/dose)
- Cefpodoxime 5 mg/kg/dose PO BID (max 200 mg/dose)
- Cefprozil 15 mg/kg/dose PO BID (max 500 mg/dose)
- Ceftriaxone 50 mg/kg/dose IM/IV qDay x 1-3 days* (daily max 1 gram/dose)
*Administer Ceftriaxone for 1 day when used as a first line for patients with penicillin allergy, and 3 days if the patient has failed other antibiotics
- Clindamycin 10 mg/kg/dose PO TID (max 600 mg/dose)

Antibiotic duration for amoxicillin, amoxicillin/clavulanate, cefuroxime, cefdinir, cefpodoxime, cefprozil, and clindamycin:

<2 years of age **OR** severe AOM **OR** chronic AOM **OR** recurrent AOM **OR** TM perforation = **10 days**
 2-5 years of age with non-severe symptoms = **7 days**
 ≥6 years of age with non-severe symptoms = **5-7 day**