




QR code for mobile view

- Inclusion Criteria:**
- Idiopathic scoliosis patients
- Exclusion Criteria:**
- Neuromuscular patients

**Inpatient (5 Henson/Hall) to Discharge**

- Key ERAS Principles:**
- Keep patient/family/team focused on early discharge
  - Advance diet, minimize IV fluids
  - Multimodal analgesia: minimize opioids, transition to orals quickly
  - Encourage time out of bed
  - Remove invasive lines (E.g., Foley)

- PCA Orders & APS Consult**
- **Hydromorphone PCA** (provide demand only if patient received methadone or IT morphine)
    - Start in PACU & to be discontinued on POD 1
  - **Ketorolac IV** 0.5 mg/kg (Max 15 mg) q6 hrs scheduled
    - Alternate with acetaminophen q3 hrs
    - Transition to PO ibuprofen 10 mg/kg (Max 800 mg) q6 hrs on POD 1
  - **Acetaminophen IV** 12.5 mg/kg (Max 750 mg) q6 hrs
    - Transition to PO acetaminophen 12.5 mg/kg (Max 750 mg) q6 hrs on POD 1
  - **Diazepam IV or PO** 0.05-0.1 mg/kg (Max 5 mg) q 4-6 hrs PRN or scheduled
  - If poor pain trajectory anticipated or if pain escalation is required, may consider addition of the following:
    - Low dose ketamine infusion
    - Dexmedetomidine infusion or clonidine IV dosing followed by patch placement
    - Other pain adjuncts as needed
  - Surgery to order **Dexamethasone IV** 0.1 mg/kg (Max 8 mg) q8 hrs x 3 doses



- Accelerated Pain Pathway**
- \*Goal to transition to PO Pain Meds on POD 0\**
- **No PCA**
  - **Ketorolac IV** 0.5 mg/kg (Max 15 mg) q6 hrs scheduled for 3 doses
    - Alternate with acetaminophen q3 hrs
    - Transition to PO ibuprofen 10 mg/kg (Max 800 mg) q6 hrs on POD 1
  - **Acetaminophen IV** 12.5 mg/kg (Max 750 mg) q6 hrs
    - Transition to PO acetaminophen 12.5 mg/kg (Max 750 mg) q6 hrs on POD 1
  - **Oxycodone PO** 0.1 mg/kg (Max 7.5 mg) q4 hrs prn
  - **Hydromorphone IV** 5 mcg/kg (Max 500 mcg) q3 hrs for breakthrough pain or not tolerating PO
- OR**
- **Morphine IV** 0.05 mg/kg (Max 4 mg) q 2 hrs prn for breakthrough pain or not tolerating PO
  - **Diazepam IV or PO** 0.05-0.1 mg/kg (Max 5 mg) q 4-6 hrs prn
  - **Dexamethasone IV** 0.1 mg/kg (Max 8 mg) q8 hrs x 3 doses

- Physical Activity**
- Ambulation**
- Encourage out of bed to chair on evening of surgery
- Physical Therapy**
- Consult started on POD 1

- Perioperative Antibiotics:**
- 1st choice: cefazolin 30 mg/kg q8 hrs X 2 doses
  - 2nd choice: clindamycin 10 mg/kg q8 hrs X 2 doses
  - MRSA: Vancomycin 15 mg/kg q12 hrs

- Diet**
- **Encourage PO intake** - advance as tolerated
  - **Initiate bowel regimen**
    - Docusate/Senna QHS on POD 0
    - Miralax BID on morning of POD 1
    - Famotidine PO 0.5 mg/kg BID
  - **Ondansetron prn for nausea/vomiting**
    - 0.1 mg/kg/dose (Max 4 mg)

- Lines, Labs, & Vitals**
- Foley Catheter**
- Remove at 0700 on POD 1
- Vital Signs**
- Vitals: q1 hrs X 4, q2 hrs for 24 hrs, q8 hrs after
  - Motor: q1 hrs X 4, q2 hrs X 4 hrs (x4), q8 hrs after
  - Neurovascular: q2 hrs X 4, q4 hrs after
- Labs**
- No routine labs scheduled

- Discharge Readiness**
- \*Discharge Goal POD2 vs POD3\**
- Discharge Requirements**
- Stable respiratory status with no oxygen requirement
  - Tolerating oral intake
  - Transitioned to oral pain medication with good pain control
  - Ambulation without assistance and cleared by PT criteria
- Discharge Teaching**
- Post-op care instructions reviewed by team with family

**Discharge home**  
 Follow-up appointments scheduled with surgeon 6 weeks postop

- [Prior to surgery algorithm](#)
- [Intraoperative care algorithm](#)