




Asthma Exacerbation: Ambulatory Algorithm



* These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.


Asthma Exacerbation: Urgent Care Algorithm


* These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.

Asthma Exacerbation: Emergency Department Algorithm


* These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.

Asthma Exacerbation: Inpatient Algorithm



* These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.

Table of Contents

Asthma Care Continuum Algorithm	1
Asthma Exacerbation: Ambulatory Algorithm	2
Asthma Exacerbation: Urgent Care Algorithm	3
Asthma Exacerbation: Emergency Department Algorithm	4
Asthma Exacerbation: Inpatient Algorithm	5
Objective of Guideline	7
Background	7
Target Users	7
Target Population	7
Practice Recommendations	8
Additional Questions Posed by the CPG Committee	9
Measures	9
Potential Cost Implications	9
Potential Organizational Barriers and Facilitators	10
Power Plans	10
Associated Policies	10
Guideline Preparation	10
Asthma Exacerbation CPG Committee Members and Representation	10
Additional Review & Feedback	11
Implementation & Follow-Up	11
Guideline Development Funding	11
Approval Process	11
Approval Obtained	11
Version History	12
References	13
Appendix A: Severity of Asthma Exacerbation	15
Appendix B: Dexamethasone versus prednisolone critically appraised topic	16
Appendix C: Albuterol dosage based on weight versus age critically appraised topic	54
Appendix D: Power Plan for Ambulatory Clinics	58
Appendix E: Quick Notes for Urgent Care	59
Appendix F: Power Plan for Emergency Department	60
Appendix G: Power Plan for Pediatric Intensive Care	61
Appendix H: Power Plan for Hospital Medicine	63
Appendix I: AGREE II Assessment for Children's Mercy Hospital's Asthma CPG	66

** These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.*

Objective of Guideline

To provide care standards for the patient diagnosed with asthma exacerbation throughout the care continuum.

Background

Efficient and effective treatment of asthma exacerbation is key to decreasing need for hospitalization, decreasing length of stay when hospitalization is required, reducing readmissions, and mitigating adverse safety events. At Children's Mercy Hospital, patients with asthma exacerbations may receive care in the ambulatory clinics, Urgent Care Clinics (UCC), Emergency Departments (ED), Medical/Surgical inpatient units, or Pediatric Intensive Care. It is imperative that we provide consistency of care and safe transitions between care settings. This Clinical Practice Guideline (CPG) provides evidence-based strategies and decision support for providers caring for patients with asthma exacerbation.

Target Users

- Physicians (Ambulatory, Urgent Care, Emergency Department, Hospital Medicine, Community Physicians, Fellows, Resident Physicians)
- Nurse Practitioners
- Nurses
- Respiratory Therapists

Target Population

Guideline Inclusion Criteria

- Patients experiencing asthma exacerbations.
- Sign and symptoms: acute onset of wheezing, coughing, and/or breathlessness with known or suspected asthma.

Guideline Exclusion Criteria

- Patients less than two years of age.
- Patients with other chronic pulmonary conditions aside from asthma.
- Long-term care of asthma without current exacerbation

AGREE

The EPR-4 national guideline and the GINA international guideline provided guidance to the Asthma Exacerbation Committee (Asthma, 2021; Expert Panel Working Group of the National Heart et al., 2020). See Tables 1 and 2 for AGREE II.

Table 1.

AGREE II^a Summary for the EPR-4 Guideline

Domain	Percent Agreement	Percent Justification
Scope and purpose	94%	The clinical questions posed, and target populations were identified. The aim of the guideline was not found in the guideline.
Stakeholder involvement	92%	The guideline was developed by the appropriate stakeholders and convened focus groups of patients and caregivers to garner input on their preferences and values. The guideline did not explicitly identify the target users, but it seems aimed at pulmonologists, allergists and PCPs.
Rigor of development	82%	The process used to gather and synthesize the evidence and the methods to formulate the recommendations were explicitly stated. The guideline developers did not provide how the guidelines will be updated.
Clarity and presentation	94%	The guideline recommendations are clear, unambiguous, and easily identified; in addition, different management options are presented.
Applicability	45%	Implementation guidance, including equipment costs and medication efficacy, were provided in the guideline. The guideline did not address barriers and facilitators that

* These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.

could be faced during implementation, monitoring or audit criteria, nor other resource costs associated with guideline implementation.

Editorial independence	94%	The recommendations were not biased with competing interests.
------------------------	-----	--

Note: Four EBP Scholars completed the AGREE II on this guideline.

Table 2.

AGREE II^a Summary for the GINA Guideline (Asthma, 2021)

Domain	Percent Agreement	Percent Justification
Scope and purpose	94%	The aim of the guideline, the clinical questions posed, and target populations were identified.
Stakeholder involvement	61%	It is unclear if the guideline included appropriate stakeholders. It is unclear if the patient's viewpoint was sought.
Rigor of development	73%	The guideline developers did not provide how the evidence was gathered and synthesized, how the recommendations were formulated.
Clarity and presentation	97%	The guideline recommendations are clear, unambiguous, and easily identified; in addition, different management options are presented.
Applicability	90%	Recommendations for monitoring adherence and treatment response are included.
Editorial independence	63%	It is unclear if the recommendations were biased by competing interests.

Note: Four EBP Scholars completed the AGREE II on this guideline.

Practice Recommendations

The National Asthma Education and Prevention Program's Expert Panel Report-3 (EPR-3) defines asthma exacerbation as an episode of "progressively worsening shortness of breathing, cough, wheezing, and chest tightness-or some combination of these symptoms" (National Asthma & Prevention, 2007). Managing asthma exacerbation in the primary or acute care settings first requires assessment of exacerbation severity based on respiratory rate, oxygen saturation, auscultation, and dyspnea (Global Initiative for Asthma, 2019; Kelly et al., 2000). Therapy with short acting beta agonist (e.g., albuterol) and supplemental oxygen, if needed, should be initiated early while assessing severity, see Appendix A (Kelly et al.) and considering alternative diagnoses (e.g., anaphylaxis; foreign body aspiration). Chest radiography and laboratory studies are not routinely needed (Global Initiative for Asthma, 2019; National Asthma & Prevention, 2007).

For severe exacerbations, immediate transfer to an acute care facility should be arranged. Intensive care may be needed for patients with lethargy, confusion, or minimal breath sounds on auscultation. Patients with severe exacerbations should be given albuterol, ipratropium bromide, magnesium sulfate, systemic corticosteroid (IV), and supplemental oxygen without delay (Global Initiative for Asthma, 2019; National Asthma & Prevention, 2007).

For moderate exacerbations, albuterol should be provided via continuous nebulization or repeated doses via metered dose inhaler (MDI) and spacer set up. Systemic corticosteroid (oral) should be given early in the course of treatment. Response to treatment should be assessed frequently to guide subsequent therapeutic interventions and assess the need for transfer to a higher level of care which may include hospitalization (Global Initiative for Asthma, 2019; National Asthma & Prevention, 2007).

For mild exacerbations, albuterol should be provided via MDI and repeated as necessary. If more than 2 doses of albuterol are required, systemic corticosteroids (oral) should be administered. Response to treatment should be assessed frequently to guide subsequent therapeutic interventions and timing of potential discharge home.

For all patients experiencing asthma exacerbations, long-term home asthma care must be addressed. Patients should be instructed to start or step-up controller therapy. Patient/family education is essential. Education should include a written asthma action plan, instructions on correct inhaler technique with emphasis on the importance of medication compliance, strategies to mitigate environmental triggers, and review of early signs of worsening asthma. Follow-up

** These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.*

within 2-7 days should be arranged (Expert Panel Working Group of the National Heart et al., 2020; Global Initiative for Asthma, 2019; National Asthma & Prevention, 2007)

Additional Questions Posed by the CPG Committee

The Expert Panel Report – 3 (EPR-3), Expert Panel Report – 4 (EPR-4), and the Global Initiative for Asthma (GINA) guidelines provided guidance to the Asthma Exacerbation Clinical Practice Guideline Committee (see Table 1 and 2 for AGREE II). While Children's Mercy adopted most of these practice recommendations, two additional questions posed by the CPG Committee led to further clarifications in care:

1. In a child > 2 years old with an acute asthma exacerbation, are 1 to 2 doses of dexamethasone as effective as 5-day course of prednisolone in the prevention of symptom recurrence?

While the Asthma CPG Committee recommends use of systemic steroid in non-intensive care settings at Children's Mercy, the committee is unable to recommend for or against the use of a one-to-two-day course of dexamethasone (intervention) in comparison to prednisolone (comparator), based on the GRADE Evidence to Decision instrument^a found in the Summary of Findings Table (see Table 1)^a. The overall certainty in the evidence is low to very low^a. Two systematic reviews and five single studies support use of dexamethasone and prednisolone in treatment of acute asthma exacerbations and both systemic steroids are effective in prevention of symptom recurrence.

*The Asthma CPG Subcommittee discussed additional considerations using the GRADE Evidence to Decision instrument^a found in the Appendix. The CPG Committee through consensus agreed on a **conditional recommendation** for dexamethasone as the systemic steroid of choice in non-intensive care settings at Children's Mercy based on feasibility, value, and compliance for all stakeholders (see Appendix B).*

2. In children aged 0 – 18 years with asthma and admitted to the hospital for an exacerbation, should the dosage of quick relief albuterol medicine via MDI **be based on weight** versus **based on age** better for improved outcomes (decreased length of stay and respiratory scores) and fewer side effects (increased HR, hyperactive, nausea/vomiting, arrhythmia, irritably)?

No recommendation can be made for weight or age-based MDI albuterol administration, based on expert review of current literature by the Department of EBP. No studies were found that answered the specific care question of weight versus age dosing for albuterol. When there is a lack of scientific evidence, standard work should be developed, implemented, and monitored (see Appendix C). To maintain consistency throughout the continuum of care at Children's Mercy, the committee opted for weight-based dosing.

Measures

- Use of Asthma Exacerbation Power Plan (UCC, ED, inpatient)
- Provision of dexamethasone as systemic steroid of choice for mild to moderate asthma exacerbations (UCC, ED, inpatient)
- Length of stay (inpatient)
- Readmissions within 72 hours of inpatient discharge
- Revisits to the UCC or ED within 72 hours within UCC or ED visit

Potential Cost Implications

The following potential improvements may reduce costs and resource utilization for healthcare facilities and reduce healthcare costs and non-monetary costs (e.g., missed school/work, loss of wages, stress) for patients and families.

- Decreased frequency of admission
- Decreased inpatient length of stay
- Decrease in readmission or acute care facility re-evaluation in less than seven days of initial exacerbation
- Decreased time to treatment in the ED setting
- Increased safety of patient transfer between settings
- Decreased unwarranted variation in care
- Narrowing gaps in health care disparities related to inequities in transportation, health literacy, and medication compliance

** These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.*

Potential Organizational Barriers and Facilitators

Barriers

- Variability of acceptable level of risk among providers
- Different clinical perspectives among providers are various care settings (acute care, subspecialty care)
- Challenges with follow-up faced by some families

Facilitators

- Collaborative engagement across care continuum settings during CPG development
- High rate of use of CPG and order sets
- Standardized order set for Urgent Care, Emergency Department, Hospital Medicine, and Pediatric Intensive Care

Power Plans

- Ambulatory Clinics (see Appendix D)
- Urgent Care (see Appendix E)
- Emergency Department (see Appendix F)
- Pediatric Intensive Care (see Appendix G)
- Hospital Medicine (see Appendix H)

Associated Policies

- Division of Emergency Medicine: Asthma Initiation Standing Order
- Continuous Albuterol Administration

Guideline Preparation

This guideline was prepared by the Evidence Based Practice (EBP) Department in collaboration with the Asthma Exacerbation CPG Committee composed of content experts at Children's Mercy Kansas City. Development of this guideline supports the Division of Service and Performance Excellence's initiative to promote care standardization that builds a culture of quality and safety that is evidenced by measured outcomes. If a conflict of interest is identified, the conflict will be disclosed next to the team member's name.

Asthma Exacerbation CPG Committee Members and Representation

- Jade Tam-Williams, MD | Pulmonology | Committee Chair
- Madison Buchanan, BHS, RRT-NPS | Respiratory Care | Committee Member
- Marc Sycip, MD | Emergency Medicine | Committee Member
- Matthew Johnson, MD | Hospital Medicine | Committee Member
- Nathan Carman, BA RRT-NPS | Respiratory Care | Committee Member
- Claire Seguin, MD | Hospital Medicine Fellow | Committee Member
- Erin Scott, DO | Emergency Medicine | Committee Member
- Aarti Pandya, MD | Allergy & Immunology | Committee Member
- Amanda Nedved, MD | Urgent Care | Committee Member
- Helen Murphy, MHS, HCEd, RRT, AE-C | Respiratory Care | Committee Member
- Caroline Holton, MD | Critical Care Fellow | Committee Member

MIT Committee Members

- George Abraham, MD | Emergency Medicine, Medical Informatics
- Ashly Catalino | Medical Informatics - Ambulatory
- Tammy Frank, RPh, CPHIMS | Medical Informatics - Pharmacy
- Brandan Kennedy, MD | Hospital Medicine, Human Factors Collaborative, Medical Informatics
- Amber Lanning | Medical Informatics - general inpatient
- Ryan McDonough, DO | Endocrinology, Medical Informatics
- Tracy Taylor | Medical Informatics - ED, UCC

** These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.*

EBP Department Members:

- Kathleen Berg, MD, FAAP | Evidence Based Practice & Hospital Medicine
- Jacqueline Bartlett, PhD, RN | Evidence Based Practice
- Andrea Melanson, OTD, OTR/L | Evidence Based Practice

Additional Review & Feedback

- The CPG was presented to each division or department represented on the CPG committee as well as other appropriate stakeholders. Feedback was incorporated into the final product.
- The CPG was reviewed by an internal and external reviewer using the AGREE II instrument (see Appendix I).

Implementation & Follow-Up

- Order sets consistent with CPG recommendations were created for each care setting (Emergency Department, Inpatient, Intensive Care).
- "Quick Orders" were updated for Urgent Care and Emergency Department.
- The Asthma Initiation Standing Order policy was updated. This details a process for nursing staff in the Emergency Department to determine severity of asthma exacerbation based on the Pediatric Asthma Score and provide albuterol and/or systemic steroids based on a standing order. This was approved by the Medical Executive Committee, Nursing Practice Council, and Pharmacy & Therapeutics Committee.
- The Continuous Albuterol Administration policy was updated to use weight-based rather than age-based albuterol dosing in all care settings in which continuous albuterol is administered. This was approved by the Pharmacy & Therapeutics Committee.
- The Respiratory Care Albuterol Weaning Protocol was updated to further standardize the dose and interval of albuterol throughout the weaning process and to maintain consistency with the CPG. This was approved by the Department of Respiratory Care.
- Education was provided to all stakeholders:
 - Nursing units where the Asthma Initiation Standing Order is used
 - Department of Respiratory Care
 - Providers from Urgent Care, Emergency Medicine, Hospital Medicine
 - Resident physicians
 Additional institution-wide announcements were made via email, hospital website, and relevant huddles.
- Metrics will be assessed and shared with appropriate care teams to determine if changes need to occur.

Guideline Development Funding

The development of this guideline was underwritten by the following departments/divisions: EBP, Pulmonology, Respiratory Care, Emergency Medicine, Hospital Medicine, Urgent Care, and Allergy & Immunology.

Approval Process

This guideline was reviewed and approved by the Asthma CPG Committee, Content Expert Departments/Divisions, and the EBP Department; after which they were approved by the Medical Executive Committee. Guidelines are reviewed and updated as necessary every 3 years within the EBP Department at CMKC. Content expert teams are involved with every review and update.

Approval Obtained

Department/Unit	Date Approved
Pulmonology	April 2022
Allergy & Immunology	April 2022
Respiratory Care	April 2022
Pediatric Intensive Care Unit	April 2022
Emergency Medicine	April 2022
Hospital Medicine	April 2022
Urgent Care	April 2022
Medical Executive Committee	July 2022

** These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.*

Version History

Date	Comments
10/2016	Version 1a: Inpatient care standards based on EPR-3 and GINA guidelines.
5/2019	Version 1b: Emergency Department and Urgent Care Clinics care standards based on EPR-3 and GINA guidelines
7/6/2022	Version two: Updated all previous guidelines (Urgent Care Clinics, Emergency Department, and Inpatient) and developed new guidelines (Care Continuum, and Ambulatory) using the EBP-4 (2020) and GINA (2021) guidelines as foundational guidelines.

Date for Next Review: July 2025

Disclaimer

When evidence is lacking or inconclusive, options in care are provided in the guideline and the power plans that accompany the guideline.

These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time.

It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.

** These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.*

References

- Abaya, R., Delgado, E. M., Scarfone, R. J., Reardon, A. M., Rodio, B., Simpkins, D., . . . Zorc, J. J. (2019). Improving efficiency of pediatric emergency asthma treatment by using metered dose inhaler. *J Asthma*, *56*(10), 1079-1086. doi:10.1080/02770903.2018.1514629
- Alangari, A. A., Malhis, N., Mubasher, M., Al-Ghamedi, N., Al-Tannir, M., Riaz, M., Umetsu, D. T., & Al-Tamimi, S. (2014). Budesonide nebulization added to systemic prednisolone in the treatment of acute asthma in children: a double-blind, randomized, controlled trial. *Chest*, *145*(4), 772-778. <https://doi.org/10.1378/chest.13-2298>
- Altamimi, S., Robertson, G., Jastaniah, W., Davey, A., Dehghani, N., Chen, R., et al. (2006). Single-dose oral dexamethasone in the emergency management of children with exacerbations of mild to moderate asthma. *Pediatric Emergency Care*, *22*(12), 786-793.
- Asthma, G. I. f. (2021). *Global Strategy for Asthma Management and Prevention*. <https://ginasthma.org/gina-reports/>
- Al-Wahadneh, A., Jboor, S., Dahabreh, M. (2006). A comparison of efficacy and safety of a single dose of intramuscular dexamethasone acetate with oral prednisolone in the management of asthma exacerbations in children. *Journal of the Royal Medical Services*, *13*(1), 15-18.
- Battistini, A. (2000). Medical treatment of respiratory emergencies. *Pediatr Med Chir*, *21*(4), 171-179.
- Children's Mercy Kansas City. (2016). *Asthma Inpatient Clinical Practice Guideline*. https://www.childrensmercy.org/contentassets/df156766f2c34b9ab72ee3eee321a71c/asthma_in_final-2016.pdf.
- Cloutier, M. M., Baptist, A. P., Blake, K. V., Brooks, E. G., Bryant-Stephens, T., DiMango, E., . . . Immunology, C. (2020). 2020 focused updates to the asthma management guidelines: a report from the National Asthma Education and Prevention Program Coordinating Committee Expert Panel Working Group. *146*(6), 1217-1270.
- Cronin, J., McCoy, S., Nally, S., Kennedy, U., Crispino-O'Connell, G., Walsh, S., (2012). A randomised trial of dexamethasone versus prednisolone in the treatment of acute paediatric asthma exacerbations. *Archives of Disease in Childhood*, *97*, A109.
- D'Vaz, N., Okitika, T. A., Shackleton, C., Devadason, S. G., & Hall, G. L. (2019). Bronchodilator responsiveness in children with asthma is not influenced by spacer device selection. *Pediatr Pulmonol*, *54*(5), 531-536. doi:10.1002/ppul.24263
- Einarsdottir, M. J., Ekman, P., Trimpou, P., Olsson, D. S., Johannsson, G., & Ragnarsson, O. (2020). High prescription rate of oral glucocorticoids in children and adults: A retrospective cohort study from Western Sweden. *Clin Endocrinol (Oxf)*, *92*(1), 21-28. <https://doi.org/10.1111/cen.14114>
- Expert Panel Working Group of the National Heart, Lung, and Blood Institute administered and coordinated National Asthma Education and Prevention Program Coordinating Committee, Cloutier, M. M., Baptist, A. P., Blake, K. V., Brooks, E. G., Bryant-Stephens, T., DiMango, E., Dixon, A. E., Elward, K. S., Hartert, T., Krishnan, J. A., Lemanske, R. F., Jr., Ouellette, D. R., Pace, W. D., Schatz, M., Skolnik, N. S., . . . Walsh, C. G. (2020). 2020 Focused Updates to the Asthma Management Guidelines: A Report from the National Asthma Education and Prevention Program Coordinating Committee Expert Panel Working Group. *J Allergy Clin Immunol*, *146*(6), 1217-1270. <https://doi.org/10.1016/j.jaci.2020.10.003>
- Fuhlbrigge, A., Peden, D., Apter, A. J., Boushey, H. A., Camargo, C. A., Jr., Gern, J., Heymann, P. W., Martinez, F. D., Mauger, D., Teague, W. G., & Blaisdell, C. (2012). Asthma outcomes: exacerbations. *J Allergy Clin Immunol*, *129*(3 Suppl), S34-48. <https://doi.org/10.1016/j.jaci.2011.12.983>
- Global Initiative for Asthma. (2019). *Global Strategy for Asthma Management and Prevention*. <https://ginasthma.org/wp-content/uploads/2019/06/GINA-2019-main-report-June-2019-wms.pdf>
- GINA. (2021). *Global Strategy of Asthma Management and Prevention*.
- Kelly, C. S., Andersen, C. L., Pestian, J. P., Wenger, A. D., Finch, A. B., Strobe, G. L., & Luckstead, E. F. (2000). Improved outcomes for hospitalized asthmatic children using a clinical pathway. *Ann Allergy Asthma Immunol*, *84*(5), 509-516. [https://doi.org/10.1016/S1081-1206\(10\)62514-8](https://doi.org/10.1016/S1081-1206(10)62514-8)
- Kirkland, S. W., Cross, E., Campbell, S., Villa-Roel, C., & Rowe, B. H. (2018). Intramuscular versus oral corticosteroids to reduce relapses following discharge from the emergency department for acute asthma. *Cochrane Database Syst Rev*, *6*(6), Cd012629. <https://doi.org/10.1002/14651858.CD012629.pub2>
- Muchão, F. P., Souza, J. M., Torres, H. C., De Lalibera, I. B., de Souza, A. V., Rodrigues, J. C., . . . da Silva Filho, L. V. (2016). Albuterol via metered-dose inhaler in children: Lower doses are effective, and higher doses are safe. *Pediatr Pulmonol*, *51*(11), 1122-1130. doi:10.1002/ppul.23469
- National Asthma Education & Prevention Program (2007). Expert Panel Report 3 (EPR-3): Guidelines for the Diagnosis and Management of Asthma-Summary Report 2007. *J Allergy Clin Immunol*, *120*(5 Suppl), S94-138. <https://doi.org/10.1016/j.jaci.2007.09.043>
- Normansell, R., Kew, K. M., & Mansour, G. (2016). Different oral corticosteroid regimens for acute asthma. *Cochrane Database Syst Rev*(5), Cd011801. <https://doi.org/10.1002/14651858.CD011801.pub2>

* These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.

- Parlar-Chun, R., & Arnold, K. (2021). Association of various weight-based doses of continuous albuterol on hospital length of stay. *J Asthma, 58*(5), 645-650. doi:10.1080/02770903.2020.1723622
- Polat, T. B., Saz, E. U., & Nursoy, M. A. (2011). Effects of salbutamol given by metered-dose inhaler on dispersion of ventricular repolarization. *Anadolu Kardiyol Derg, 11*(3), 232-236. doi:10.5152/akd.2011.063
- Ratnayake, A., Taveras, H., Iverson, H., & Shore, P. (2016). Pharmacokinetics and pharmacodynamics of albuterol multidose dry powder inhaler and albuterol hydrofluoroalkane in children with asthma. *Allergy Asthma Proc, 37*(5), 370-375. doi:10.2500/aap.2016.37.3985
- Schuh, S., Johnson, D. W., Stephens, D., Callahan, S., Winders, P., & Canny, G. J. (1999). Comparison of albuterol delivered by a metered dose inhaler with spacer versus a nebulizer in children with mild acute asthma. *J Pediatr, 135*(1), 22-27. doi:10.1016/s0022-3476(99)70322-7
- Schuh, S., Zemek, R., Plint, A., Black, K. J., Freedman, S., Porter, R., . . . Johnson, D. W. (2012). Practice patterns in asthma discharge pharmacotherapy in pediatric emergency departments: a pediatric emergency research Canada study. *Acad Emerg Med, 19*(9), E1019-1026. doi:10.1111/j.1553-2712.2012.01433.x

* These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.

Appendix A: Severity of Asthma Exacerbation

Severity of Asthma Exacerbation			
	Mild	Moderate	Severe
Respiratory Rate	2-3 yrs: ≤ 34 /min 4-5 yrs: ≤ 30 /min 6-12 yrs: ≤ 26 /min >12 yrs: ≤ 23 /min	2-3 yrs: 35-39/min 4-5 yrs: 31-35/min 6-12 yrs: 27-30/min >12 yrs: 24-27/min	2-3 yrs: ≥ 40 /min 4-5 yrs: ≥ 36 /min 6-12 yrs: ≥ 31 /min >12 yrs: ≥ 28 /min
Pulse Oximetry (SpO₂) Requirement	>95% on room air	90-95% on room air	<90% on room air
Auscultation	Normal or end expiratory wheeze	Expiratory wheezing	Inspiratory and expiratory wheeze or diminished breath sounds
Retractions	None or intercostal	Intercostal and substernal	Intercostal, substernal, and supraclavicular
Dyspnea	Speaks in sentences	Speaks in partial sentences	Speaks in single words or short phrases
<i>Patients may not display all signs/symptoms in a single column. Clinical judgement is essential to assessment of severity.</i>			

** These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interest of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.*

Appendix B: Dexamethasone versus prednisolone critically appraised topic

Specific Care Question #1:

In a child greater than 2 years old with an acute asthma exacerbation, are 1-2 doses of dexamethasone (intervention) as effective as a 5-day course of prednisolone (comparator) in prevention of symptom recurrence?

Recommendations from the Asthma CPG Committee and Based on Current Literature

While the Asthma Clinical Practice Guideline (CPG) Committee recommends use of systemic steroid in non-intensive care settings at Children's Mercy, the committee is unable to recommend for or against the use of a one-to-two-day course of dexamethasone (intervention) in comparison to prednisolone (comparator), based on the GRADE Evidence to Decision instrument^a found in the Summary of Findings Table (see Table 1)^a. The overall certainty in the evidence is low to very low^a. Two systematic reviews and five single studies support use of dexamethasone and prednisolone in treatment of acute asthma exacerbations and both systemic steroids are effective in prevention of symptom recurrence.

The Asthma CPG Subcommittee discussed additional considerations using the GRADE Evidence to Decision instrument^a found in the Appendix. The CPG Committee through consensus agreed on a conditional recommendation for dexamethasone as the systemic steroid of choice in non-intensive care settings at Children's Mercy based on feasibility, value, and compliance for all stakeholders (see Appendix).

Literature Summary

Background

Acute asthma exacerbations are a leading cause for patients seeking emergent medical care at acute care centers and, although most patients are discharged within the same day, relapse of symptoms is still common requiring additional medical care and return to an acute care center (Kirkland et al., 2018). Systemic corticosteroids are a primary part of the treatment regimen for moderate to severe asthma exacerbations with dexamethasone and prednisolone most often prescribed (Fuhlbrigge et al., 2012). In spite of the proven efficacy of dexamethasone and prednisolone, these steroids, along with others, require the balance of benefits against the potential adverse events such as nausea, vomiting, or gastrointestinal distress (Normansell et al., 2016). Evidence is limited to which medications and dosing provide maximum recovery from acute exacerbations in children, specifically to decrease relapse in symptoms. This review will summarize identified literature to answer the specific care question.

Study Characteristics

The search for suitable studies was completed on September 8, 2021. Amanda Nedved, MD, Erin Scott, DO, and Claire Seguin, MD reviewed the 42 titles and/or abstracts found in the search and identified^b five systematic reviews and six single studies believed to answer the question. After an in-depth review of the identified systematic reviews^a and single studies^a, two systematic reviews and five single studies answered the question.

Race/Ethnicity Race and ethnicity as defined by the individual authors were reviewed in the literature. Of the three studies that reported on race and ethnicity, 50-70% of participants were either black or Hispanic.

Are one to two doses of dexamethasone as effective as a five-day course of prednisolone in prevention of symptom recurrence?

Elkharwili et al. (2020) recruited 60 patients aged 2-11 years and randomized into three groups. For purposes of this review, only group 1: single dose of 0.3 mg/kg dexamethasone and group 3: five days of 1.5 mg/kg/day prednisolone were compared for relapse rate of symptoms over five days.

Hermani et al. (2021) completed a retrospective review of 1,410 patients aged 3-21 years of age. The authors measured relapse of symptoms based on two interventions: receipt of dexamethasone or prednisolone prior to presentation to the emergency department (ED) and receipt of

dexamethasone or prednisolone after ED presentation. All four groups received oral dexamethasone (0.5 mg/kg/day for a median of 1 day) or prednisolone (average dose of 1.8 mg/kg/day for median of 2 days) after arrival to the hospital.

Kirkland et al. (2018), a systematic review, reported on both adult and pediatric studies to analyze the optimal delivery method (oral or intramuscular) of dexamethasone compared to oral prednisolone. Only the pediatric studies are included in this review (Al-Wahadneh et al., 2006; Gordon et al., 2007; Gries et al., 2000; Klig et al., 1997). The primary outcome of relapse of symptoms was defined by the authors as any unscheduled visit to a health practitioner for worsening asthma symptoms or requiring subsequent treatment with corticosteroids. Reported relapse data within 10 days of discharge from the ED were reported.

Normansell et al. (2016), a systematic review, reviewed both adult and pediatric studies to analyze higher dose/longer course versus lower dose/shorter course for the outcome of re-admission during the follow-up period. Only the pediatric studies are included in this review (Altamimi et al., 2006; Cronin et al., 2015; Greenberg et al., 2008; Qureshi et al., 2001). The pediatric studies compared a single dose (0.6 mg/kg) of dexamethasone to a five-day dosing of prednisolone (2mg/kg). Relapse of symptoms, up to 15 days post discharge from the ED, was used as the parameters for follow up.

Paniagua et al. (2017) analyzed data on 557 randomized patients aged 1-14 years comparing the impact of two doses of dexamethasone to five days of prednisolone for relapse of symptoms defined as a return visit to the ED.

Volk et al. (2019) completed a retrospective review of a two-day course of dexamethasone to a five-day course of prednisolone on symptom recurrence within one week of initial visit to a hospital emergency department.

Watnick et al. (2016) analyzed the impact of a single dose of dexamethasone to a three-to-five-day course of prednisolone on relapse of symptoms in patients presenting to an area emergency room aged 3-17 years. Those that returned within 72 hours of discharge from the emergency room were counted as having a relapse but were only counted for their initial return.

Summary by Outcome

Relapse of Symptoms with 1 Day of Dexamethasone vs. 3-5 Days of Prednisolone.

Four studies (Elkharwili et al., 2020; Kirkland et al., 2018; Normansell et al., 2016; Watnick et al., 2016) measured the relapse in symptoms of an asthma exacerbation within 14 days following initial presentation, ($n = 9,424$). Based on the pediatric studies ($n = 615$) identified in the two systematic reviews (Kirkland et al., 2018; Normansell et al., 2016), the $OR = 0.74$, 95% CI [0.32, 1.69], $p = .47$ indicated the intervention of one day dosing of dexamethasone was not different to the comparator of three to five days dosing of prednisolone (see Figure 2 & Table 1). For the RCT study (Elkharwili et al., 2020), ($n = 8,769$), the $OR = 0.63$, 95% CI [.40, 1.01], $p = .05$ indicated the intervention of one day dosing of dexamethasone was not different to the three to five days dosing of prednisolone (see Figure 3 & Table 1). The cohort study (Watnick et al., 2016), ($n = 40$), $MD = 3.00$, 95% CI [-14.67, 20.67], $p = .74$ indicated the intervention of one day dosing of dexamethasone was not different to the comparator of three to five days dosing of prednisolone (see Figure 4 & Table 1).

Certainty of the Evidence For Relapse of Symptoms with 1 Day of Dexamethasone vs. 3-5 Days of Prednisolone. The certainty of the body of evidence was low to very low. The body of evidence for the two systematic reviews (Kirkland et al., 2018; Normansell et al., 2016) was assessed to have serious risk of bias as demonstrated by lack of blinding of study participants and study personnel and serious imprecision due to low number of events ($n = 35$). The body of evidence for the RCT (Elkharwili et al., 2020) was found to have serious risk of bias as demonstrated by data analysis completed per protocol and very serious imprecision as demonstrated by a low number of subjects ($n = 40$). The one retrospective cohort study (Watnick et al., 2016) was assessed to have very serious imprecision as demonstrated by low number of events ($n = 164$).

Relapse of Symptoms with 2 Days of Dexamethasone vs. 5-6 Days of Prednisolone.

Three studies (Normansell et al., 2016; Paniagua et al., 2017; Volk et al., 2019) measured the relapse in symptoms of an asthma exacerbation within 14 days of the initial exacerbation, ($n = 1,342$). For the one systematic review (Normansell et al., 2016), using two of the pediatric studies (Greenberg et al., 2008; Qureshi et al., 2001) and the single RCT (Paniagua et al., 2017) met the criteria for review, ($n = 1,279$), the $OR = 1.65$, 95% CI [.85, 3.19], $p = .14$, indicated the intervention of two day dosing of dexamethasone was not different to the comparator of five to six day dosing of prednisolone (see Figure 5 & Table 2). The one cohort study (Volk et al., 2019), ($n = 63$), the $OR = .33$, 95% CI [.02, 7.13], $p = .48$, indicated the intervention of two-day dosing of dexamethasone was not different to five-to-six-day dosing of prednisolone (see Figure 6 & Table 2).

Certainty of the Evidence for Relapse of Symptoms with 2 Days of Dexamethasone vs. 5-6 Days of Prednisolone. The certainty of the body of evidence was low for the one systematic review and one RCT but very low for the observational study. The body of evidence for the systematic review (Normansell et al., 2016) and the RCT (Paniagua et al., 2017), was assessed to have serious risk of bias due to study participants and study personnel not blinded causing concern for performance bias. The observational study (Volk et al., 2019) was assessed to have very serious imprecision due to small number of events and subjects.

Relapse of Symptoms with 2 Doses of Dexamethasone vs. 5 Days of Prednisolone initiated after hospital arrival.

One study (Hermani et al., 2021) measured the relapse in symptoms of an asthma exacerbation within 10 days of the initial exacerbation, ($n = 961$). For the outcome of relapse of symptoms, the $OR = 6.20$, 95% CI [0.37, 103.50], $p = .20$ indicated the intervention of two days of dexamethasone was not different compared to five days of prednisolone initiated after hospital arrival (see Table 3).

Certainty of the Evidence for Relapse of Symptoms with 2 Doses of Dexamethasone vs. 5 days of Prednisolone initiated after hospital arrival. The certainty of the body of evidence was very low. The body of evidence for the one observational study (Hermani et al., 2021) was assessed to have serious imprecision due to a low number of events and subjects. As only one study (Hermani et al., 2021) was identified to answer this question inconsistency could not be assessed.

Relapse of Symptoms with 1-3 doses of Dexamethasone vs. 1-3 doses of Prednisolone before hospital arrival.

One study (Hermani et al., 2021) measured the relapse in symptoms of an asthma exacerbation within 10 days of the initial exacerbation, ($n = 449$). For the outcome of relapse of symptoms, the $OR = .76$ 95% CI [.14, 3.94], $p = .74$ indicated the intervention of one to three doses of dexamethasone was not different than one to three doses of prednisolone provided prior to hospital arrival in decreasing relapse of asthma symptoms (see Table 4).

Certainty of the Evidence for Relapse of Symptoms with 1-3 doses of Dexamethasone vs. 1-3 doses of Prednisolone before hospital arrival. The certainty of the body of evidence was very low. The body of evidence for the one observational study (Hermani et al., 2021) was assessed to imprecision due to low number of events. As only one study (Hermani et al. 2021) was identified to answer this question, inconsistency could not be assessed.

Identification of Studies

Search Strategy and Results (see Figure 1)

"Status Asthmaticus"[Mesh] OR "Asthma/drug therapy"[Mesh] OR "asthma exacerbation*" AND ("Dexamethasone/administration and dosage"[Mesh] OR "Prednisolone/administration and dosage"[Mesh] OR "Prednisone/administration and dosage"[Mesh]) AND (child OR children OR pediatr* OR paediatr* OR infant OR adolescence
 Records identified through database searching $n = 41$
 Additional records identified through other sources $n = 1$

Studies Included in this Review

Citation	Study Type
----------	------------

*Elkharwili et al., 2020	RCT
Hermani et al., 2021	Cohort
*Kirkland et al., 2018	SR
* Normansell et al., 2016	SR
*Paniagua et al., 2017	RCT
Volk et al., 2019	Cohort
Watnick et al., 2016	Cohort

References marked with an asterisk indicate studies included in the meta-analysis

Studies Not Included in this Review with Exclusion Rationale

Citation	Reason for exclusion
SR Bravo-Soto et al., 2017	In Spanish language
SR Kirkland et al., 2019	Articles of interest are included in two of the included SR
SR Meyer et al., 2014	Articles of interest are included in in Kirkland et al. (2018) SR

Methods Used for Appraisal and Synthesis

^aThe GRADEpro Guideline Development Tool (GDT) is the tool used to create the Summary of Findings table(s) for this analysis.

^bRayyan is a web-based software used for the initial screening of titles and / or abstracts for this analysis (Ouzzani, Hammady, Fedorowicz & Elmagarmid, 2017).

^cReview Manager (Higgins & Green, 2011) is a Cochrane Collaborative computer program used to assess the study characteristics as well as the risk of bias and create the forest plots found in this analysis.

^dThe Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram depicts the process in which literature is searched, screened, and eligibility criteria is applied (Moher, Liberati, Tetzlaff, & Altman, 2009).

References to Appraisal and Synthesis Methods

^aGRADEpro GDT: GRADEpro Guideline Development Tool (2015). McMaster University, (developed by Evidence Prime, Inc.). [Software]. Available from grade.pro.org.

^bOuzzani, M., Hammady, H., Fedorowicz, Z., & Elmagarmid, A. (2016). Rayyan-a web and mobile app for systematic reviews. *Systematic Reviews*, 5(1), 210. doi:10.1186/s13643-016-0384-4

^cHiggins, J. P. T., & Green, S. e. (2011). *Cochrane Handbook for Systematic Reviews of Interventions [updated March 2011]* (Version 5.1.0 ed.): The Cochrane Collaboration, 2011.

^dMoher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 6(7): e1000097. doi:10.1371/journal.pmed1000097 **For more information, visit www.prisma-statement.org.**

Question Originator

A.Nedved, MD, E. Scott, DO, and C. Seguin, MD

Medical Librarian Responsible for the Search Strategy

K. Swaggart, MLIS, AHIP

EBP Team or EBP Scholar’s Responsible for Analyzing the Literature

J. A. Bartlett, PhD, RN

J. Dusin, MS, RD, LD, CPHQ

B. Hunter, RN, BSN, CPN

J. Wierson, RN, BSN, MBA, CCRC

EBP Medical Director Responsible for Reviewing this Document

K. Berg, MD, FAAP

EBP Team Member Responsible for Reviewing, Synthesizing, and Developing this Document

A. Melanson, OTD, OTR/L


Acronyms Used in this Document

Acronym	Explanation
AGREE II	Appraisal of Guidelines Research and Evaluation II
CAT	Critically Appraised Topic
EBP	Evidence Based Practice
ED	Emergency Department
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analyses

Statistical Acronyms Used in this Document

Statistical Acronym	Explanation
CI	Confidence Interval
I^2	Heterogeneity test
M or \bar{X}	Mean
n	Number of cases in a subsample
N	Total number in sample
OR	Odds Ratio
P or p	Probability of success in a binary trial
RCT	Randomized controlled trial
RR	Relative risk
SD	Standard deviation
SE	Standard error
SR	Systematic Review

Figure 1



Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)^d

Summary of Findings Table
Table 1

Summary of Findings Table^a: Relapse of Symptoms 1 Day Dexamethasone vs. 3-5 Days Prednisolone

Certainty assessment							Summary of findings					
Participants (studies) Follow-up	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias	Overall certainty of evidence	Study event rates (%)		Relative effect (95% CI)	Anticipated absolute effects		
							With 5-day course of prednisolone	With 1-2 doses of dexamethasone		Risk with 5-day course of prednisolone	Risk difference with 1-2 doses of dexamethasone	
Relapse of symptoms (1 day Dexamethasone vs. 3-5 days Prednisolone)												
615 (6 RCTs)	serious ^a	not serious	not serious	serious ^b	none	⊕⊕○○ Low	19/299 (6.4%)	16/316 (5.1%)	OR 0.74 (0.32 to 1.69)	64 per 1,000	16 fewer per 1,000 (from 42 fewer to 39 more)	
Relapse of symptoms (1 day Dexamethasone vs. 3-5 days Prednisolone)												
40 (1 RCT)	serious ^d	not serious	not serious	very serious ^c	none	⊕○○○ Very low	20	20	MD = 3.00 (-14.67, 20.67)	The mean relapse of symptoms (1 day Dexamethasone vs. 3-5 days Prednisolone) was 0	MD 3 higher (14.67 lower to 20.67 higher)	
Relapse of symptoms (1 day Dexamethasone vs. 3-5 days Prednisolone)												
8769 (1 observational study)	not serious	not serious	not serious	serious ^b	none	⊕⊕○○ Low	143/7130 (2.0%)	21/1639 (1.3%)	OR 0.63 (0.40 to 1.01)	20 per 1,000	7 fewer per 1,000 (from 12 fewer to 0 fewer)	

- Notes:
- a. both study participants and study personnel not blinded, concerns for performance bias
 - b. low number of events
 - c. low number of subjects
 - d. randomization not completed as stated and data analysis followed per protocol analysis

Table 2
Summary of Findings Table^a: Relapse of symptoms 2 days Dexamethasone vs. 5-6 days Prednisolone

Certainty assessment							Summary of findings				
Participants (studies) Follow-up	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias	Overall certainty of evidence	Study event rates (%)		Relative effect (95% CI)	Anticipated absolute effects	
							With 5-day course of prednisolone	With 1-2 doses of dexamethasone		Risk with 5-day course of prednisolone	Risk difference with 1-2 doses of dexamethasone
Relapse of symptoms (2 days Dexamethasone vs. 5-6 days Prednisolone)											
1279 (3 RCTs)	serious ^a	not serious	not serious	serious ^b	none	⊕⊕○○ Low	15/675 (2.2%)	25/604 (4.1%)	OR 1.65 (0.85 to 3.19)	22 per 1,000	14 more per 1,000 (from 3 fewer to 45 more)
Relapse of symptoms (2 days Dexamethasone vs. 5-6 days Prednisolone)											
63 (1 observational study)	not serious	not serious	not serious	serious ^c	none	⊕○○○ Very low	2/40 (5.0%)	0/23 (0.0%)	OR 0.33 (0.02 to 7.13)	50 per 1,000	33 fewer per 1,000 (from 49 fewer to 223 more)

- Notes:
- a. both study participants and study personnel not blinded, concerns for performance bias
 - b. low number of events
 - c. Low number of events and subjects

Table 3
Summary of Findings Table^a: Relapse of symptoms 2 days Dexamethasone vs. 5 days Prednisolone initiated after hospital arrival hospitalized

Certainty assessment							Summary of findings				
Participants (studies) Follow-up	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias	Overall certainty of evidence	Study event rates (%)		Relative effect (95% CI)	Anticipated absolute effects	
							With 5-day course of prednisolone	With 1-2 doses of dexamethasone		Risk with 5-day course of prednisolone	Risk difference with 1-2 doses of dexamethasone
Relapse of symptoms (2 doses Dexamethasone vs. 5 doses Prednisolone during hospitalization)											
961 (1 observational study)	not serious	not serious	not serious	serious ^a	none	⊕○○○ Very low	0/135 (0.0%)	18/826 (2.2%)	OR 6.20 (0.37 to 103.50)	0 per 1,000	0 fewer per 1,000 (from 0 fewer to 0 fewer)

- Notes:
- If you have questions regarding this Specific Care Question – please contact evidencebasedpractice@cmh.edu

a. Low number of events and subjects

Table 4
Summary of Findings Table^a: Relapse of symptoms 1-3 doses Dexamethasone vs. 1-3 doses Prednisolone before hospital arrival

Certainty assessment							Summary of findings				
Participants (studies) Follow-up	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias	Overall certainty of evidence	Study event rates (%)		Relative effect (95% CI)	Anticipated absolute effects	
							With 5-day course of prednisolone	With 1-2 doses of dexamethasone		Risk with 5-day course of prednisolone	Risk difference with 1-2 doses of dexamethasone
Relapse of symptoms (1-3 doses Dexamethasone vs. 1-3 doses Prednisolone before hospital arrival)											
449 (1 observational study)	not serious	not serious	not serious	serious ^a	none	⊕○○○ Very low	5/294 (1.7%)	2/155 (1.3%)	OR 0.76 (0.14 to 3.94)	17 per 1,000	4 fewer per 1,000 (from 15 fewer to 47 more)

Notes:

a. Low number of events and subjects

Meta-analysis

Figure 2

Comparison: 1 day Dexamethasone versus 3-5 days Prednisolone, Outcome: Relapse of symptoms





Figure 3

Comparison: 1 day Dexamethasone versus 3-5 days Prednisolone, Outcome: Relapse of symptoms



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Figure 4
Comparison: 1 day Dexamethasone versus 3-5 days Prednisolone, Outcome: Relapse of symptoms






Figure 5
Comparison: 2-day Dexamethasone versus 5-6 days Prednisolone, Outcome: Relapse of symptoms



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Figure 6
Comparison: 2-day Dexamethasone versus 5-6 days Prednisolone, Outcome: Relapse of symptoms



Characteristics of Intervention Studies
Elkharwili, 2020

Methods	Randomized Control Trial
<p>Participants</p>	<p>Participants: Children with acute exacerbation of asthma Setting: Hospital (Tanta University Hospital, Egypt, March 2016 - October 2017) Randomized into study: <i>N</i> = 94</p> <ul style="list-style-type: none"> • Group 1, 0.3 mg/kg oral dexamethasone for one day: <i>n</i> = 35 • Group 2, 0.6 mg/kg of oral dexamethasone for two days: <i>n</i> = 32 • Group 3, 1.5 mg/kg oral prednisolone: <i>n</i> = 27 <p>Completed Study Treatment: <i>N</i> = 81</p> <ul style="list-style-type: none"> • Group 1: <i>n</i> = 29 • Group 2: <i>n</i> = 29 • Group 3: <i>n</i> = 23 <p>Completed Follow-up Phase of Study: <i>N</i> = 60</p> <ul style="list-style-type: none"> • Group 1: <i>n</i> = 20 • Group 2: <i>n</i> = 20 • Group 3: <i>n</i> = 20 <p>Gender, males (as defined by researchers):</p> <ul style="list-style-type: none"> • Group 1: <i>n</i> = 40% • Group 2: <i>n</i> = 50% • Group 3: <i>n</i> = 55% <p>Race / ethnicity or nationality:</p> <ul style="list-style-type: none"> • The authors did not identify race or ethnicity of the participants. <p>Age, mean in years (SD):</p> <ul style="list-style-type: none"> • Group 1: 5.93 (2.37) • Group 2: 6.52 (2.64) • Group 3: 6.15 (2.75) <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> • children with a history of bronchial asthma, • those that presented with an asthma exacerbation, which was defined as a decrease in expiratory airflow that could be documented and quantified by simple measurement of lung function (spirometry or peak expiratory flow (PEF)) • age 2 - 11 years

	<ul style="list-style-type: none"> • male or female <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> • children aged 11 years • children with intubation history for previous asthma exacerbations • children with active varicella or herpes simplex infection in the past 3 weeks • children with documented concurrent infection with respiratory syncytial virus • use of oral or intravenous corticosteroids in the previous 4 weeks • concurrent stridor • known patients with tuberculosis • presence of other significant comorbidities such as: cardiac, immune, liver, endocrine, neurological and psychiatric disorders <p>Power Analysis: Analysis at a p value of 0.05 and a power of 80% showed that a total sample size of 78 patients distributed as 1:1:1 in the three groups was necessary. The level of significance was set at a p value < 0.05, while p values of 0.01 and 0.001 were considered highly significant.</p>
<p>Interventions</p>	<p>Group 1: single dose of 0.3 mg/kg oral dexamethasone, with a maximum dose of 12mg/day for 1 day and continued with a placebo for the other 4 days</p> <p>Group 2: 0.6 mg/kg of oral dexamethasone, with a maximum dose of 16 mg/day in three divided doses for two consecutive days and continued with a placebo for the other 3 days</p> <p>Group 3: 1.5 mg/kg oral prednisolone per day for 5 days with a maximum dose of 60 mg in three divided doses</p>
<p>Outcomes</p>	<p>Primary outcome(s):</p> <ul style="list-style-type: none"> • Change in physical examination, Pediatric Respiratory Assessment Measure (PRAM) score*, the Modified Pulmonary Index Score (MPIS)*, pulmonary function tests*, saturated oxygen, blood eosinophilic count and serum immunoglobulin E after 5 days of taking the corticosteroids <p>Secondary outcome:</p> <ul style="list-style-type: none"> • Vomiting, gastrointestinal tract (GIT) cramps and relapse rate were recorded as secondary outcomes of the study <p>Safety Outcome:</p> <ul style="list-style-type: none"> • Relapse Rate* <p>*Outcomes of interest for the CPG Team</p>

<p>Notes</p>	<ul style="list-style-type: none"> • Due to protocol deviations and participants lost to follow-up, the authors did not meet the sample size calculated to determine significance • There was no statistically significant difference in weight gain and blood sugar before and after 5 days of treatment within the same group • After 5 days of treatment, pairwise comparison showed a significant difference in blood sugar level only between group II and group III (p=0.004) • After 5 days of treatment, comparison of the participants showed that there was a highly statistical difference in MPIS, oxygen need, duration of hospital admission and PRAM within the three groups (p<0.001). • After 30 days, ATAZ Asthma Therapy Assessment Questionnaire (ATAQ) showed no significant differences among the three studied groups for missed days of school
---------------------	--

<i>Risk of bias table</i>		
Bias	Judgement	Support for judgement
Random sequence generation (selection bias)	High risk	Article states that 94 eligible patients were assigned and randomized in a 1:1:1 ratio into three groups. It does not specify as to how the randomization was generated. Although this is stated, it shows that the following were the initial group allocations: Group I: 35 patients, Group II: 32 patients and Group III: 27 patients which does not prove that a 1:1:1 ratio was used.
Allocation concealment (selection bias)	Unclear risk	Insufficient information to permit judgment of low or high risk
Blinding of participants and personnel (performance bias)	Unclear risk	Article states that it was a double-blind clinical trial but doesn't describe any further information regarding blinding methods
Blinding of outcome assessment (detection bias)	Unclear risk	Insufficient information to permit judgment of low risk or high risk
Incomplete outcome data (attrition bias)	High risk	The authors identify in Table 1 that patients with protocol deviations (Group I: 6, Group II: 3, Group III: 4) were not counted as completing study. In Table 5 the authors only include in the final analysis the data from only the participants completing the follow-up phase therefore data is missing from 21 additional participants (Group I: 9, Group II: 9 and Group III: 3). With the removal of this data the authors did not meet the sample size needed to detect significance between the different groups.
Selective reporting (reporting bias)	Low risk	The only thing not noted in the outcomes table was the saturation oxygen, but there were other parameters captured such as PEF (%) and FEV1/FVC (%) so noted as low risk
Other bias	Unclear risk	There may be a risk of bias, but there is insufficient information to assess whether an important risk of bias exists.

Hemani, 2021

Methods	Multisite Retrospective Cohort
<p>Participants</p>	<p>Participants: Patients 3 to 21 years admitted between January 1, 2013, and December 31, 2017, with primary discharge diagnosis, IDC 9 and ICD 10, of asthma exacerbation or status asthmaticus Setting: Atlanta, USA, Tertiary Children's Hospital System Number enrolled into study: <i>N</i> = 1410</p> <ul style="list-style-type: none"> • Group 1, Dexamethasone (DEX) Initiated After Hospital Arrival: <i>n</i> = 826 • Group 2, Prednisone/prednisolone (PRED) Initiated After Hospital Arrival: <i>n</i> = 135 • Group 3, Dexamethasone (DEX) Before Hospital Arrival: <i>n</i> = 155 • Group 4, Prednisone/prednisolone (PRED) Before Hospital Arrival: <i>n</i> = 294 <p>Gender, males:</p> <ul style="list-style-type: none"> • Group 1: <i>n</i> = 531 (64.3%) • Group 2: <i>n</i> = 77 (57%) • Group 3: <i>n</i> = 96 (62%) • Group 4: <i>n</i> = 192 (65.3%) <p>Race (reported numbers do not reach total enrolled, but reported percentages equal 100):</p> <ul style="list-style-type: none"> • Black <ul style="list-style-type: none"> ○ Group 1: <i>n</i> = 562 (72.3%) ○ Group 2: <i>n</i> = 76 (58%) ○ Group 3: <i>n</i> = 83 (55%) ○ Group 4: <i>n</i> = 152 (53.3%) • White <ul style="list-style-type: none"> ○ Group 1: <i>n</i> = 126 (16.2%) ○ Group 2: <i>n</i> = 33 (35%) ○ Group 3: <i>n</i> = 43 (29%) ○ Group 4: <i>n</i> = 76 (26.7%) • Asian <ul style="list-style-type: none"> ○ Group 1: <i>n</i> = 17 (2.2%) ○ Group 2: <i>n</i> = 1 (1%) ○ Group 3: <i>n</i> = 6 (4%) ○ Group 4: <i>n</i> = 8 (2.8%) • Other <ul style="list-style-type: none"> ○ Group 1: <i>n</i> = 72 (9.3%) ○ Group 2: <i>n</i> = 20 (15%) ○ Group 3: <i>n</i> = 18 (12%) ○ Group 4: <i>n</i> = 49 (17.2%)

Ethnicity:

- Hispanic or Latino
 - **Group 1:** $n = 111$ (13.5%)
 - **Group 2:** $n = 26$ (19%)
 - **Group 3:** $n = 19$ (12%)
 - **Group 4:** $n = 31$ (10.6%)
- Non- Hispanic or Latino
 - **Group 1:** $n = 714$ (86.6%)
 - **Group 2:** $n = 109$ (81%)
 - **Group 3:** $n = 136$ (88%)
 - **Group 4:** $n = 262$ (89.4%)

Age, mean in years, (SD):

- **Group 1:** 6.79 (3.3)
- **Group 2:** 6.54 (3.1)
- **Group 3:** 6.49 (3.3)
- **Group 4:** 6.87 (3.1)

Inclusion Criteria:

- Age of 3 to 21 years
- Receiving monotherapy with DEX or PRED
- Multiple asthma-related hospital visits within a 10-day period only the first encounter was captured

Exclusion Criteria:

- Less than 3 years of age
- Receiving an unspecified oral steroid or combination of DEX and PRED during acute illness
- Missing information about steroid administration prior to admission
- Methyl prednisone administration during acute illness
- Steroid administration in the prior 2 weeks or receiving a prolonged steroid course
- Initial PICU admission
- Concurrent diagnosis of bronchiolitis, pneumonia, or croup
- Use of Bi-level positive airway pressure
- Supplemental therapies in the Emergency Department (e.g., antibiotics, oseltamivir, heliox, terbutaline, racemic epinephrine, hypertonic saline, chest physiotherapy, and budesonide)
- Pulmonary or cardiac comorbidities, sickle cell disease, down syndrome, or immunosuppression
- Hospital admissions with paper chart documentation
- Left against medical advice or readmission

	<p>Covariates Identified:</p> <ul style="list-style-type: none"> • Albuterol administration prior to hospital arrival
<p>Interventions</p>	<p>Both: Received a clinical respiratory score; received albuterol; may receive ipratropium, magnesium, and supplemental oxygen</p> <ul style="list-style-type: none"> • Group 1: Received an average dose of DEX 0.5 mg/kg per day for a median of 2 days while hospitalized • Group 2: Received an average dose of PRED 1.8 mg/kg per day for a median of 2 days while hospitalized • Group 3: Received an average dose of DEX 0.5 mg/kg per day for a median of 1 day while hospitalized • Group 4: Received an average dose of PRED 1.8 mg/kg per day for a median of 2 days while hospitalized
<p>Outcomes</p>	<p>Primary outcome:</p> <ul style="list-style-type: none"> • Length of stay (LOS)* <p>Secondary outcomes:</p> <ul style="list-style-type: none"> • PICU transfer during initial hospitalization* • Readmission within 10 days after hospital discharge* <p>Safety outcome:</p> <ul style="list-style-type: none"> • Not reported <p>*Outcomes of interest to the CMH CPG /CAT development team</p>
<p>Notes</p>	<p>Limitations:</p> <ul style="list-style-type: none"> • Retrospective study, susceptible to adjustment items • Majority of patients classified as mild intermittent or mild persistent asthma • Exclusion criteria prevented severe asthma exacerbation patient inclusion in study • Previous inhaled corticosteroid uses not included • Steroid adherence after discharge not tracked

Kirkland, 2018

<p>Methods</p>	<p>Systematic Review (meta-analysis)</p>
<p>Objective</p>	<p>To examine the effectiveness and safety of a single dose of intramuscular (IM) corticosteroids provided prior to discharge compared to a short course of oral corticosteroids in the treatment of acute asthma patients discharged from an ED or equivalent acute care setting.</p>
<p>Methods</p>	<p>Criteria for considering studies for this review</p> <ul style="list-style-type: none"> • Types of studies: RCTs or controlled clinical trials • Participants: Adults and pediatric patients presenting with acute asthma to an ED or acute care setting. • Target Condition(s): Acute asthma exacerbation <p>Search methods for identification of studies</p> <ul style="list-style-type: none"> • Electronic databases searched: Cochrane Airways Group Register of Trials and databases including Medline, Embase, EBM ALL, Global Health, International Pharmaceutical Abstracts, CINAHL, SCOPUS, ProQuest Dissertations and Theses Global, and LILACS. • Search strategy employed: <ul style="list-style-type: none"> ○ *Secondary Prevention; Acute Disease; Administration, Oral; Adrenal Cortex Hormones [*administration & dosage]; Anti-Asthmatic Agents [*administration & dosage]; Asthma [drug therapy] [*prevention & control]; Betamethasone [administration & dosage]; Dexamethasone [administration & dosage]; Emergency Service, Hospital; Injections, Intramuscular; Methylprednisolone [administration & dosage]; Patient Discharge; Prednisolone [administration & dosage]; Prednisone [administration & dosage]; Randomized Controlled Trials as Topic; Recurrence; Triamcinolone [administration & dosage] ○ Included full text, abstracts, and unpublished data in search criteria. • Searching other resources (such as reference list): Reference lists of all primary studies and review articles were checked for additional references. Relevant manufacturers' web sites were also searched for additional study information. <p>Data collection and analysis</p> <ul style="list-style-type: none"> • Inclusion criteria: <ul style="list-style-type: none"> ○ RCTs and controlled clinical trials ○ Studies with acute exacerbation of asthma as primary reason for presentation to ED with no other co-existing complications ○ Asthma diagnosis had to be made either using international/national clinical guidelines or spirometric criteria or both • Exclusion criteria: <ul style="list-style-type: none"> ○ Studies that focused on corticosteroid treatment in hospitalized patients • Population: Adult and pediatric patients with uncomplicated exacerbation of asthma • Setting: Hospital ED or equivalent acute care setting • Study Design: Systematic review and meta-analysis • Data collection process: Two independent reviewers assessed study eligibility and study quality. Disagreements were resolved by a third party and assessed the risk of bias using the Cochrane Risk of Bias toll.

	<ul style="list-style-type: none"> • Assessment of the certainty of the evidence: Quality of the evidence was measured/assessed using GRADE. • Data Synthesis (what statistical plan do the authors establish a priori): <ul style="list-style-type: none"> ○ Random effects model used and performed a sensitivity analysis with a fixed-effect model. ○ Heterogeneity: I^2 statistic used to measure heterogeneity. If substantial heterogeneity was identified, it was reported, and possible causes were explored using a prespecified subgroup analysis (see subgroup analysis below): <ul style="list-style-type: none"> ▪ Children (zero to 18 years of age) versus adults (18 years of age and older) to examine any potential age-specific treatment effects of IM or oral corticosteroids. ▪ Relapse occurring within 10 days and over 10 days post-discharge. ▪ Low versus moderate versus high exacerbation severity based on the pulmonary function taken at the time of the participant's presentation to the ED, prior to treatment with a bronchodilator. ▪ Co-interventions received (ICS versus ICS corticosteroids/long-acting beta₂-agonists (LABA)). ○ Sensitivity analysis carried out by removing the following types of studies from primary outcome analyses: <ul style="list-style-type: none"> ▪ Studies that we consider to be at high risk of bias based on the criteria outlined in the Cochrane Handbook for Systematic Reviews of Interventions (Higgins 2011). ▪ Studies in which the duration of oral corticosteroid treatment was less than five days. ▪ The results from fixed-effect models were compared with the random-effects models for the main outcome. ▪ Studies in which supplemental corticosteroids were provided to the patients in the ED as a co-intervention
<p>Results</p>	<p>Study Selection (actual results/data) Number of articles identified: $N = 912$ Full-text articles assessed for eligibility: $n = 20$</p> <ul style="list-style-type: none"> ○ Studies included in qualitative synthesis: $n = 9$ <p>Synthesis of quality of evidence (strength of evidence): Using GRADE, the overall certainty of the evidence was assessed per outcome ranging from low to moderate with the following results per outcome:</p> <ul style="list-style-type: none"> • Primary outcomes of relapse as well as relapse after 10 days was rated as low quality due to overall unclear to high risk of bias of the studies and imprecision due to wide confidence intervals including both benefit, harm and no effect. The subgroup analysis for relapse was rated at low quality due to the low number of available patients and wide confidence intervals. • Outcome for adverse events also ranked at low quality due to overall unclear to high risk of bias of the included studies and imprecision due to few events. • Outcome of symptom persistence and 24-hour beta₂-agonists use ranked as low quality due to the overall unclear to high risk of bias of the included studies as well as few events. • Outcome of peak expiratory flow ranked as moderate quality due to imprecision of the results.

	<p>Synthesis of quantitative evidence: (pediatric patients only)</p> <ul style="list-style-type: none"> • Overall Effect Size: Intramuscular versus oral corticosteroids, Outcome: Relapse <ul style="list-style-type: none"> ▪ Odds Ratio: .78 ▪ CI: 95% CI [.38, 1.57], <i>p</i> = .49 ▪ Heterogeneity <ul style="list-style-type: none"> • <i>I</i>²=0% • Overall Effect Size: Intramuscular versus oral corticosteroids, Outcome: Relapse intention to treat <ul style="list-style-type: none"> ▪ Odds Ratio: .78 ▪ CI: 95% CI [0.38, 1.57], <i>p</i> = .48 ▪ Heterogeneity <ul style="list-style-type: none"> • <i>I</i>²=0% • Overall Effect Size: Intramuscular versus oral corticosteroids, Outcome: Relapse within 10 days <ul style="list-style-type: none"> ▪ Odds Ratio: .75 ▪ CI: 95% CI [0.28, 2.0], <i>p</i> = .57 • Overall Effect Size: Intramuscular versus oral corticosteroids, Outcome: Relapse over 10 days <ul style="list-style-type: none"> ▪ Odds Ratio: .78 ▪ CI: 95% CI [0.38, 1.57], <i>p</i> = .48 ▪ Heterogeneity <ul style="list-style-type: none"> • <i>I</i>²=0%
<p>Discussion</p>	<p>Summary of evidence</p> <ul style="list-style-type: none"> • Systemic corticosteroids were found to be an effective treatment in decreasing relapse of symptoms for acute asthma exacerbation for ED or equivalent acute care settings and assists with prevention of admission however, the optimal route of dosing and administration is unclear. <p>Limitations</p> <ul style="list-style-type: none"> • Lack of reporting out of data on secondary outcomes significantly limited the number of studies that could be used for the meta-analysis and impacted the authors’ ability to draw meaningful conclusions or recommendations towards the overall effectiveness of IM corticosteroids. • Only four pediatric studies met the inclusion criteria. • The effectiveness of the corticosteroids results may have been impacted by the age of the children enrolled in the study meaning, younger children may not respond to the corticosteroids due to fewer airway eosinophils. • Co-interventions were poorly reported in studies reviewed and it is likely that some of the agents used may no longer be used. • Lack of reporting on the use of the ICS and ICS/LABA agents limited the review on its ability to estimate the impact of these agents on the efficacy of IM or oral corticosteroids. • Dosing of corticosteroids was not a criterion used for inclusion and thus, no conclusion was drawn on the impact of dosing completed.
<p>Funding</p>	<p>The National Institute for Health Research (NIHR) supported this project, via Cochrane Infrastructure funding to the Cochrane Airways Group.</p>

Normansell, 2016

Methods	Systematic Review (meta-analysis)
Objective	To assess the efficacy and safety of any dose or duration of oral steroids versus any other dose or duration of oral steroids for adults and children with asthma exacerbation.
Methods	<p>Criteria for considering studies for this review</p> <ul style="list-style-type: none"> • Types of studies: <ul style="list-style-type: none"> ○ RCTs • Participants: <ul style="list-style-type: none"> ○ Adults ○ Children • Target Condition(s): <ul style="list-style-type: none"> ○ Acute Asthma Attack <p>Search methods for identification of studies</p> <ul style="list-style-type: none"> • Electronic databases searched: the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE, EMBASE, the Cumulative Index to Nursing and Allied Health Literature (CINAHL), the Allied and Complementary Medicine Database (Alangari et al.) and PsycINFO, and by handsearching respiratory journals and meeting abstracts • Search strategy employed: Mesh terms (see study for full list) • Searching other resources: Handsearching of respiratory journals and meeting abstracts <p>Data collection and analysis</p> <ul style="list-style-type: none"> • Inclusion criteria: <ul style="list-style-type: none"> ○ Randomized controlled trials (RCTs), irrespective of blinding or duration, that evaluated one dose or duration of oral steroid versus any other dose or duration, for management of asthma exacerbations. ○ Both adults and children with asthma of any severity, in which investigators analyzed adults and children separately. ○ Other co-intervention in the management of an asthma exacerbation, provided it was not part of the randomized treatment. • Exclusion criteria: <ul style="list-style-type: none"> ○ Wrong comparator ○ Wrong intervention ○ Not randomized • Population: <ul style="list-style-type: none"> ○ Adults and children with acute exacerbation of asthma • Setting: <ul style="list-style-type: none"> ○ Inpatient ○ Emergency department • Study Design: <ul style="list-style-type: none"> ○ Systematic review and meta-analysis • Data collection process: <ul style="list-style-type: none"> ○ Data collection form designed by two of the investigators

	<ul style="list-style-type: none"> • Assessment of the certainty of the evidence: <ul style="list-style-type: none"> ○ GRADE • Data Synthesis (what statistical plan do the authors establish a priori): <ul style="list-style-type: none"> ○ Overall Effect Size (just state what is being used in the study) <ul style="list-style-type: none"> ▪ OR ▪ RD ▪ CI ○ Heterogeneity <ul style="list-style-type: none"> ▪ Cochran's Q statistic ▪ I² statistic
<p style="text-align: center;">Results</p>	<p>Study Selection (actual results/data) Number of articles identified: <i>N</i> = 1406 Full-text articles assessed for eligibility: <i>n</i> = 71 <ul style="list-style-type: none"> ○ Studies included in qualitative synthesis: <i>n</i> = 18 Synthesis of quality of evidence (strength of evidence): <ul style="list-style-type: none"> • Low to very low certainty Synthesis of quantitative evidence: <ul style="list-style-type: none"> • Prednisolone vs dexamethasone, outcome: Admission during follow-up <ul style="list-style-type: none"> ○ OR = .09 (-0.07, 0.26), <i>p</i>-value = .9 ○ <i>n</i> = 3 studies (985 patients) ○ I² = 0% • Prednisolone vs dexamethasone, outcome: Re-admission during follow-up <ul style="list-style-type: none"> ○ OR = .44 (0.15, 1.33), <i>p</i>-value = .14 ○ <i>n</i> = 3 studies (985 patients) ○ I² = 0% • Prednisolone vs dexamethasone, outcome: Asthma symptoms: Pulmonary Index Score <ul style="list-style-type: none"> ○ MD = -.1 (0.45, 0.25), <i>p</i>-value = .58 ○ <i>n</i> = 1 study (100 patients) • Prednisolone vs dexamethasone, outcome: Asthma symptoms: Patient Self-Assessment Score <ul style="list-style-type: none"> ○ MD = .1 (-0.67, 0.69), <i>p</i>-value = .98 ○ <i>n</i> = 1 study (100 patients) • Prednisolone vs dexamethasone, outcome: Asthma symptoms: Pediatric Respiratory Assessment Measure <ul style="list-style-type: none"> ○ MD = 0 (-0.36, 0.36), ○ <i>n</i> = 1 study (218 patients) • Prednisolone vs dexamethasone, outcome: new exacerbation during follow-up period: unscheduled visit to healthcare provider <ul style="list-style-type: none"> ○ OR = .85 (0.54, 1.34), <i>p</i>-value = .48 ○ <i>n</i> = 4 study (981 patients) ○ I² = 0% </p>

	<ul style="list-style-type: none"> • Prednisolone vs dexamethasone, outcome: new exacerbation during follow-up period: oral corticosteroids prescribed <ul style="list-style-type: none"> ○ OR = .29 (0.1, 0.81) ○ <i>n</i> = 1 study (242 patients)
<p>Discussion</p>	<p>Summary of evidence</p> <ul style="list-style-type: none"> • There was difficulty combining the results of studies in a useful way because investigators used a variety of doses and durations of steroids and measured their results in diverse ways. Also, events such as hospital admissions and serious side effects happened very rarely in these studies, making it difficult to tell whether longer or shorter courses or higher or lower doses are better or safer, or if prednisolone is generally better or worse than dexamethasone. Some studies were old and did not use steroid doses or durations used by medical practitioners today. <p>Limitations</p> <ul style="list-style-type: none"> • Evidence presented in the review is generally considered to be of low or very low certainty, which means there is a great amount of uncertainty of whether the results are accurate, mostly because the authors could not combine many studies. Some studies did not clearly explain how trial organizers decided which people would receive which dose of steroids, and in some studies, both participants and trial organizers knew which dose they were getting.
<p>Funding</p>	<p>Funding</p> <ul style="list-style-type: none"> • Cochrane Collaborative

Paniagua, 2017

Methods	Randomized Control Trial
Participants	<p>Participants: Children with asthma exacerbation who presented to the emergency department (ED) Sept 2014-October 2015</p> <p>Setting: Acute care teaching tertiary hospital, Spain (Basque Country)</p> <p>Randomized into study: <i>N</i> = 590</p> <ul style="list-style-type: none"> • Group 1: Dexamethasone, <i>n</i> = 294 • Group 2: Prednisolone, <i>n</i> = 296 <p>Completed Study: <i>N</i> = 557</p> <ul style="list-style-type: none"> • Group 1: <i>n</i> = 281 • Group 2: <i>n</i> = 276 <p>Gender, males: mean, (%)</p> <ul style="list-style-type: none"> • Group 1: <i>n</i> = 169 (60.1%) • Group 2: <i>n</i> = 166 (60.1%) <p>Race / ethnicity or nationality:</p> <ul style="list-style-type: none"> • Not reported <p>Age, years (mean) (Einarsdottir et al.):</p> <ul style="list-style-type: none"> • Group 1: 4.7 (3.4) • Group 2: 4.5 (3.4) <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> • Aged 1-14 years • History of previous diagnosis of asthma or at least 2 previous episodes responsive wheeze or first wheezing episode in a child > 2 years with history of atopy • Respiratory symptoms- <ul style="list-style-type: none"> ○ Acute cough, shortness of breath, tachypnea attributed to bronchospasm (wheezing, prolong expiration), increased work of breathing, and/or increased bronchodilator requirements from baseline <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> • Other airway pathology • Other diseases that require hospitalization for safety • Children with life-threatening asthma exacerbation • Use of oral or parenteral corticosteroids in the past 4 weeks <p>Power Analysis: Sample size calculation was based on a Pediatric Asthma Control Tool (PACT) score at day seven for the dexamethasone group would not be more than 6% greater than the prednisolone group score; a sample size of at least 556 subjects was required to detect a difference.</p>

Interventions	<p>Both groups: received the first 2-3 β_2-agonist treatments at 20-minute intervals with the addition of ipratropium bromide prescribed per attending provider.</p> <ul style="list-style-type: none"> • Group 1: Dexamethasone*, oral, (1 mg/ml), 0.6 mg/kg, maximum 12 mg, one dose received in the ED, a second dose was administered 24 hours later. • Group 2: Prednisone/prednisolone*, oral, 1.5 mg/kg, maximum 60 mg, one dose in the ED, followed by 1 mg/kg/d, maximum 60 mg, twice daily on days 2 - 5. Choice of liquid or tablet formulate was based on the subject's age. <ul style="list-style-type: none"> ○ *If either treatment was vomited within 30 minutes, the dose was re-administered. <p>Subjects were contacted by phone on day 7 and 15 in which PACT questionnaire and the asthma related quality of life (ARQoL) instrument was completed. Both instruments are validated.</p>
Outcomes	<p>Primary outcome(s):</p> <ul style="list-style-type: none"> • Percent of subjects with symptoms at 7 days [PACT score] * and their quality of life score [ARQoL score]. <p>Secondary outcome(s):</p> <ul style="list-style-type: none"> • Vomiting • Adherence to treatment • Parent satisfaction • Admission rate* • Unscheduled returns to ED* • Hospital re-admissions • Visits to Primary Care Provider • School and work absenteeism <p>Safety outcome(s):</p> <ul style="list-style-type: none"> • Not reported <p>*Outcomes of interest to the CMH CPG team</p>
Notes	<p>Trial registered - clinicaltrialsregister.eu: 2013-003145-42, the registry states it is ongoing July 2, 2018,</p>

<i>Risk of bias table</i>		
Bias	Judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Statisticians performed the randomization
Allocation concealment (selection bias)	Low risk	allocation concealment was maintained using sequentially numbered opaque envelopes containing a letter A (experimental treatment) or B (conventional treatment), following the randomization list.
Blinding of participants and personnel (performance bias)	High risk	Open label, with subjective outcomes
Blinding of outcome assessment (detection bias)	Unclear risk	Data managers and the statistical team were blinded but bias could have occurred during the interview with family.

Incomplete outcome data (attrition bias)	Low risk	Used a per-protocol analysis, met sample size needed to detect inferiority between interventions.
Selective reporting (reporting bias)	Low risk	All outcomes were reported
Other bias	High risk	Treating physician was permitted to exclude patients if time constraints made enrollment unfeasible. The PACT tool used in a six-item inventory. References were found to the 10 and 3 item PACT, not the 6 item PACT. Self-reported response to both the PACT and the quality-of-life inventories.

Volk, 2019

Methods	Retrospective Cohort
<p>Participants</p>	<p>Participants: Pediatric patients with Asthma or wheezing, Setting: Ambulatory Setting between August 2013 to July 2015 Number enrolled into study: <i>N</i> = 63</p> <ul style="list-style-type: none"> • Group 1, Prednisone: <i>n</i> = 40 • Group 2, Dexamethasone: <i>n</i> = 23 <p>Gender, males (as defined by researchers):</p> <ul style="list-style-type: none"> • Group 1: <i>n</i> = 31 (78%) • Group 2: <i>n</i> = 23 (78%) <p>Race / ethnicity or nationality (as defined by researchers):</p> <ul style="list-style-type: none"> • Group 1: Non-Hispanic <i>n</i> = 16 (40%) • Group 1: Hispanic <i>n</i> = 24 (60%) • Group 2: Non-Hispanic <i>n</i> = 6 (26%) • Group 2: Hispanic <i>n</i> = 17 (74%) <p>Age, mean (years)</p> <ul style="list-style-type: none"> • Group 1: 6.4 • Group 2: 7.8 <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> • ≥ 3 years of age • Primary visit diagnosis of “wheezing” (ICD9 786.07), “asthma unspecified type with exacerbation” (ICD9 493.92), “asthma with status asthmaticus” (ICD9 493.91), or “cough variant asthma” (ICD9 493.82) <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> • Received steroid treatment from an outside health facility within 1-week of presentation to the Center <p>Covariates Identified:</p> <ul style="list-style-type: none"> • Not reported

<p>Interventions</p>	<p>Both: inhaled β-agonist treatment prior to corticosteroid with supplemental oxygen is oxygen saturations fall below 94%.</p> <ul style="list-style-type: none"> • Group 1: Oral Prednisone-a single dose of weight-based prednisolone as either an oral tablet or liquid solution. Additional daily single doses are prescribed and completed at home over 5 days. • Group 2: Oral Dexamethasone-single dose of a dissolvable oral tablet using a weight-based formula at the Center. A second dose is prescribed and given within 24 hrs. (typically at home) to complete the 2-day course
<p>Outcomes</p>	<p>Primary outcome(s):</p> <ul style="list-style-type: none"> • ED visits • Hospital admissions • Return clinic visits within 1 week for recurrent • *Persistent symptoms <p>*Outcomes of interest to the CMH CPG development team</p>
<p>Notes</p>	<p>Results:</p> <ul style="list-style-type: none"> • The rates of hospital admissions, ED visits, and symptom follow-up were similar between the 2 groups ($P > .05$). • The cost for a course of dexamethasone was US \$1.28 versus US \$16.20 for prednisolone. The average cost for an asthma exacerbation office visit was US \$79.89 compared with US \$3113.28 for an ED visit. <p>Limitations:</p> <ul style="list-style-type: none"> • As the EMR was surveyed, errors may exist in coding and documentation • Unable to determine the true illness severity as measured by the number of previous exacerbations and the dose or duration of inhaled corticosteroids • Call backs were not done to determine medication compliance or medication adverse effects • Insurance claims from outside health facilities could not be tracked for 16% of patients, do not know if they were treated for wheezing elsewhere

Watnick, 2016

Methods	Cohort
<p>Participants</p>	<p>Participants: patients 3 to 17 years old with acute asthma exacerbations Setting: urban tertiary care children's hospital ED Number enrolled into study: <i>N</i> =13,518 (4,749 excluded because they did not receive corticosteroid) number included in study: 8,769</p> <ul style="list-style-type: none"> • Group 1, prednisone/prednisolone: <i>n</i> = 7130 • Group 2, dexamethasone: <i>n</i> = 1639 <p>Gender, males (as defined by researchers)-not described per study group but overall patients compared to those with corticosteroids and relapse:</p> <ul style="list-style-type: none"> • <i>n</i> = 8,281 (61%) (all patients with & without corticosteroid treatment) • <i>n</i> = 109 (60 %) (patients with relapse) <p>Race / ethnicity or nationality (as defined by researchers):</p> <ul style="list-style-type: none"> • 4,783 (35%) White (all patients with & without corticosteroid treatment) 63 (34%) White (patients with relapse) • 7,701 (57%) Black (all patients with & without corticosteroid treatment) 108 (59%) Black (patients with relapse) • 119 (1%) Asian (all patients with & without corticosteroid treatment) 1 (1%) Asian (patients with relapse) • 36 (0%) American Indian or Alaskan (all patients with & without corticosteroid treatment) 0 (0%) American Indian or Alaskan (patients with relapse) • 1 (0%) Pacific Islander (all patients with & without corticosteroid treatment) 0 (0%) (patients with relapse) • 878 (7%) unknown or declined (all patients with & without corticosteroid treatment) 11 (6%) (patients with relapse) <p>Age, mean/median in months/years, (range/IQR)</p> <ul style="list-style-type: none"> • Group 1: 7 (4-10) (all patients with & without corticosteroid treatment) • Group 2: 7 (4-11) (patients with relapse) <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> • Patients 3 to 17 years old • Seen in ED, treated with systemic corticosteroids and subsequently discharged • Those that returned within 72 hours with continued asthma symptoms <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> • Patients in ED for asthma exacerbation not receiving corticosteroids or IV formulation of corticosteroids • For patients with multiple return trips to the ED within 72 hours, only the first return visit was analyzed. <p>Covariates Identified:</p> <ul style="list-style-type: none"> • None identified

Interventions	<ul style="list-style-type: none"> • Group 1: oral prednisone or prednisolone-2 mg/kg for 3 to 5 days • Group 2: oral dexamethasone 0.6mg/kg given in a single dose
Outcomes	<p>Primary outcome(s):</p> <ul style="list-style-type: none"> • *Relapse rates of patients receiving oral dexamethasone with those receiving oral prednisone or prednisolone. <p>Secondary outcome(s):</p> <ul style="list-style-type: none"> • None described <p>Safety outcome(s):</p> <ul style="list-style-type: none"> • None <p>*Outcomes of interest to the CMH CPG development team</p>
Notes	<p>Results:</p> <ul style="list-style-type: none"> • Group 1: 143 cases of relapse of symptoms • Group 2: 21 cases of relapse of symptoms <p>Limitations:</p> <ul style="list-style-type: none"> • Lack of information available on patient's severity of asthma exacerbation • Lack of information on detailed asthma characteristics, patient's exposure to smoke, and flu vaccine status • Potential loss of patients that would have qualified for the study inclusion, however, may have been classified incorrectly from the International Classification of Diseases, ninth edition

Evidence to Decision for Dexamethasone

Should 1-2 doses of dexamethasone vs. 5-day course of prednisolone be used for children greater than 2 years old with acute asthma exacerbation?	
POPULATION:	children greater than 2 years old with acute asthma exacerbation
INTERVENTION:	1-2 doses of dexamethasone
COMPARISON:	5-day course of prednisolone
MAIN OUTCOMES:	Relapse of symptoms (1 day Dexamethasone vs. 3-5 days Prednisolone); Relapse of symptoms (1 day Dexamethasone vs. 3-5 days Prednisolone); Relapse of symptoms (1 day Dexamethasone vs. 3-5 days Prednisolone); Relapse of symptoms (2 days Dexamethasone vs. 5-6 days Prednisolone); Relapse of symptoms (2 days Dexamethasone vs. 5-6 days Prednisolone); Relapse of symptoms (2 doses Dexamethasone vs. 5 doses Prednisolone during hospitalization); Relapse of symptoms (1-3 doses Dexamethasone vs. 1-3 doses Prednisolone before hospital arrival);

ASSESSMENT

Problem Is the problem a priority?		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
<ul style="list-style-type: none"> o No o Probably no o Probably yes ● Yes o Varies o Don't know 	<p>Since the last review of asthma exacerbations in pediatrics, there has been an uptick in literature measuring the efficacy of 1-2 doses of dexamethasone compared to a 5-day course of prednisolone. Dexamethasone is less expensive with a long half-life compared to prednisolone. In addition, prednisolone's poor palatability can make compliance with a five-day course challenging, especially with children. Thus, the question becomes a priority if providers have an alternative systemic corticosteroid that demonstrates similar recovery of symptoms yet is both less expensive and requires fewer doses.</p>	
Desirable Effects How substantial are the desirable anticipated effects?		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
<ul style="list-style-type: none"> o Trivial o Small o Moderate ● Large o Varies o Don't know 	<p>In review of all studies, the test for overall effect showed the intervention (dexamethasone) and the control (prednisolone) were effective and equivalent in reducing relapse of symptoms regardless of dosing provided.</p>	<p>The desired anticipated effect is substantial considering the consequences of relapse of symptoms. Relapse may lead to missed school/work, repeat ambulatory visits, repeat ED visits, or readmission.</p>

<p>Undesirable Effects How substantial are the undesirable anticipated effects?</p>		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
<ul style="list-style-type: none"> <input type="radio"/> Large <input checked="" type="radio"/> Moderate <input type="radio"/> Small <input type="radio"/> Trivial <input type="radio"/> Varies <input type="radio"/> Don't know 	<p>Nausea, vomiting, and GI distress are noted undesirable effects of both dexamethasone and prednisolone. Side effects (SMD 0.03; 95% CI (-0.38, 0.44) in the first 7-10 days, while rarely reported, showed no differences between the treatment groups (Rowe, B. H., Spooner, C. H., Ducharme, F. M., Bretzlaff, J. A., & Bota, G. W., 2001).</p>	<p>Theoretically, a longer treatment course may increase the risk of adrenal suppression.</p> <p>Anecdotally, the committee notes more neuropsychiatric side effects (labile mood, poor sleep) with prednisolone compared to dexamethasone.</p>
<p>Certainty of evidence What is the overall certainty of the evidence of effects?</p>		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
<ul style="list-style-type: none"> <input type="radio"/> Very low <input checked="" type="radio"/> Low <input type="radio"/> Moderate <input type="radio"/> High <input type="radio"/> No included studies 	<p>While systemic corticosteroids are standard of care for asthma exacerbation, the overall certainty of the evidence is low to very low that dexamethasone vs prednisolone show differences in relapse of symptoms.</p>	
<p>Values Is there important uncertainty about or variability in how much people value the main outcomes?</p>		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
<ul style="list-style-type: none"> <input type="radio"/> Important uncertainty or variability <input type="radio"/> Possibly important uncertainty or variability <input checked="" type="radio"/> Probably no important uncertainty or variability <input type="radio"/> No important uncertainty or variability 	<p>As there was no substantial difference with effect on relapse of symptoms for dexamethasone compared to prednisolone and the variability between studies reflects a 'no difference' in outcome, clinicians are left to determine best choice of corticosteroid for their setting based on ease of provision and likelihood of compliance. However, there is probably no important variability as to how much clinicians value the outcome of no relapse of symptoms.</p>	

<p>Balance of effects Does the balance between desirable and undesirable effects favor the intervention or the comparison?</p>		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
<ul style="list-style-type: none"> ○ Favors the comparison ○ Probably favors the comparison ○ Does not favor either the intervention or the comparison ● Probably favors the intervention ○ Favors the intervention ○ Varies ○ Don't know 	<p>No difference in desirable or undesirable effects were found to support either dexamethasone or prednisolone within the literature reviews.</p>	<p>Consideration of additional effects (other than relapse of symptoms) favors the intervention (dexamethasone). Dexamethasone is easier to administer (often 1 dose in the care setting before discharge home), less expensive, and essentially eliminates the issue of noncompliance. Noncompliance with prednisolone could be related to treatment duration, poor palatability, side effects, cost and/or the process of filling the prescription.</p>
<p>Resources required How large are the resource requirements (costs)?</p>		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
<ul style="list-style-type: none"> ○ Large costs ○ Moderate costs ○ Negligible costs and savings ● Moderate savings ○ Large savings ○ Varies ○ Don't know 	<p>Outside of CM (Children s Mercy), prednisolone costs for a five-day course can range from \$18.00 to \$48.00 compared to dexamethasone pricing for a one-to-two-day course costs \$11.00 to \$32.00 based on insurance and pharmacy.</p>	<p>Overall, dexamethasone cost for the treatment course is less than that of prednisolone.</p> <p>According to CM standard charges for 2022, the self-pay costs per unit are as follows: Dexamethasone 12mg/12ml oral solution - \$11.77 Dexamethasone 4mg tablet - \$8.29 Prednisolone 3mg/ml oral solution - \$4.16 x 5 days Prednisone 10mg tab - \$3.88 x 5 days Prednisone 20mg tab - \$3.97 x 5 days</p> <p>Additional costs include the time, effort, and transportation needed to get a prednisolone prescription filled at a pharmacy, compared to receiving dexamethasone in the care setting prior to discharge.</p>

<p>Certainty of evidence of required resources What is the certainty of the evidence of resource requirements (costs)?</p>		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
<ul style="list-style-type: none"> ○ Very low ○ Low ● Moderate ○ High ○ No included studies 	<p>The majority of patients will take either dexamethasone or the first dose of prednisolone in the care setting (urgent care, emergency department, inpatient) so cost for initial dosing would be the same regarding resources of staff and staging. The only difference would be the cost in drug pricing.</p>	
<p>Cost effectiveness Does the cost-effectiveness of the intervention favor the intervention or the comparison?</p>		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
<ul style="list-style-type: none"> ○ Favors the comparison ○ Probably favors the comparison ○ Does not favor either the intervention or the comparison ○ Probably favors the intervention ● Favors the intervention ○ Varies ○ No included studies 	<p>The cost effectiveness would favor the dexamethasone (intervention) with an average of \$7.00 to \$16.00 less, depending on insurance and pharmacy. Additional cost savings for dexamethasone include no need for time or transportation to go to a pharmacy.</p>	
<p>Equity What would be the impact on health equity?</p>		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
<ul style="list-style-type: none"> ○ Reduced ○ Probably reduced ○ Probably no impact ● Probably increased ○ Increased ○ Varies ○ Don't know 	<p>Fifty percent to 70% of participants were either of Black race or Hispanic ethnicity. The majority of initial visits were through a medical care settings' emergency department.</p> <p>The use of dexamethasone allows for equal efficacy (based on relapse of symptoms) without the impact of inequalities potentially posed by prednisolone. Some subpopulations may have more challenges related to transportation to a pharmacy and medication costs/medical insurance. Literacy or language barriers may impact the efficacy of prescription instructions.</p>	

<p>Acceptability Is the intervention acceptable to key stakeholders?</p>		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
<ul style="list-style-type: none"> <input type="radio"/> No <input type="radio"/> Probably no <input type="radio"/> Probably yes <input checked="" type="radio"/> Yes <input type="radio"/> Varies <input type="radio"/> Don't know 	<p>It is acceptable to key stakeholders to use an equally effective, yet less expensive medication. Stakeholders also value the increased ease of administration (fewer doses, better palatability) of the intervention (dexamethasone) which may improve compliance.</p>	
<p>Feasibility Is the intervention feasible to implement?</p>		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
<ul style="list-style-type: none"> <input type="radio"/> No <input type="radio"/> Probably no <input type="radio"/> Probably yes <input checked="" type="radio"/> Yes <input type="radio"/> Varies <input type="radio"/> Don't know 	<p>The intervention is feasible to implement. It is available in CM urgent care, emergency department, and inpatient settings. The first dose of systemic corticosteroid is already given in the care setting, so the use of dexamethasone does not create additional processes. Medication access and administration of dexamethasone is more feasible than prednisolone for patients and their families.</p>	

SUMMARY OF JUDGEMENTS

	JUDGEMENT						
PROBLEM	No	Probably no	Probably yes	Yes		Varies	Don't know
DESIRABLE EFFECTS	Trivial	Small	Moderate	Large		Varies	Don't know
UNDESIRABLE EFFECTS	Large	Moderate	Small	Trivial		Varies	Don't know
CERTAINTY OF EVIDENCE	Very low	Low	Moderate	High			No included studies
VALUES	Important uncertainty or variability	Possibly important uncertainty or variability	Probably no important uncertainty or variability	No important uncertainty or variability			
BALANCE OF EFFECTS	Favors the comparison	Probably favors the comparison	Does not favor either the intervention or the comparison	Probably favors the intervention	Favors the intervention	Varies	Don't know
RESOURCES REQUIRED	Large costs	Moderate costs	Negligible costs and savings	Moderate savings	Large savings	Varies	Don't know
CERTAINTY OF EVIDENCE OF REQUIRED RESOURCES	Very low	Low	Moderate	High			No included studies
COST EFFECTIVENESS	Favors the comparison	Probably favors the comparison	Does not favor either the intervention or the comparison	Probably favors the intervention	Favors the intervention	Varies	No included studies
EQUITY	Reduced	Probably reduced	Probably no impact	Probably increased	Increased	Varies	Don't know
ACCEPTABILITY	No	Probably no	Probably yes	Yes		Varies	Don't know
FEASIBILITY	No	Probably no	Probably yes	Yes		Varies	Don't know

TYPE OF RECOMMENDATION

Strong recommendation against the intervention	Conditional recommendation against the intervention	Conditional recommendation for either the intervention or the comparison	Conditional recommendation for the intervention	Strong recommendation for the intervention
○	○	○	●	○

Appendix C: Albuterol dosage based on weight versus age critically appraised topic

Specific Care Question #2

In children aged 0 – 18 years with asthma and admitted to the hospital for an exacerbation, should the dosage of quick relief albuterol medicine via metered dose inhaler (MDI) **be based on weight** versus **based on age** better for improved outcomes (decreased length of stay and respiratory scores) and fewer side effects (increased HR, hyperactive, nausea/vomiting, arrhythmia, irritably).

Recommendations Based on Current Literature (Best Evidence) Only

No recommendation can be made for weight or age-based MDI albuterol administration, based on expert review of current literature by the Department of EBP. No studies were found that answered the specific care question of weight versus age dosing for albuterol. When there is a lack of scientific evidence, standard work should be developed, implemented, and monitored.

Literature Summary

Background Asthma is a chronic disease characterized by airway inflammation (Global Initiative for Asthma (GINA), 2021). Respiratory symptoms such as chest tightness, cough, shortness of breath, wheezing, and variable expiratory airflow are common citation. Symptoms can be chronic or occur suddenly, with acute amplification of symptoms (GINA, 2021). An accepted treatment for mild-to-moderate exacerbation is administering short-acting beta agonists (SABA), such as albuterol, administered through an MDI (GINA, 2021). The previous dosing recommendations have been based on the number of puffs given through MDI (Children’s Mercy Kansas City, 2016). The purpose of this review is to determine if weight-based versus age-based dosing results in improved outcomes.

Two guidelines were identified for this review (Cloutier et al., 2020; GINA, 2021). Both guidelines were assessed using AGREE II (see Table 1).

The Global Initiative for Asthma (2020) and The National Asthma Education and Prevention Program Coordinating Committee Working Group Expert Panel Report (EPR)-4 (Cloutier et al., 2020) do not make any recommendations for short-acting beta-agonists (SABA) based on age or weight.

Medication	Dose	Comments
Albuterol MDI (90 mcg/puff)	4-8 puffs every 20 minutes for 3 doses, then every 1-4 hours inhalation maneuver as needed. Add mask in children <4 years	In mild-to-moderate exacerbation, MDI plus valved-holding chamber is as effective as nebulized therapy with appropriate administration technique and coaching by trained personnel

(Cloutier et al., 2020)

Medication	Dose	Comments
Albuterol MDI (90 mcg/puff)	4-10 puffs every 20 minutes for the first hour, After the first hour, doses vary from 4-10 puffs every 3-4 hours up to 6-10 puffs every 1-2 hours, or more often	Mild-to-moderate exacerbation, delivery of SABA via MDI and spacer leads to similar improvement in lung functions as delivery via nebulizer

(GINA, 2021)

Study Characteristics The search for suitable studies was completed on April 1, 2021. H. Murphy, BHS RRT AE-C and M. Buchanan BHS, RRT-NPS reviewed the 76 titles and/or abstracts found in the search and identified^a two guidelines and nine single studies believed to answer the question. After an in-depth review of the identified guidelines and single studies, none answered the specific care question, but one guideline addressed provided general recommendations related to the question.

Identification of Studies

Search Strategy and Results (see Figure 1)

((“Asthma”[Majr]) AND “Metered Dose Inhalers”[Mesh]) AND “Albuterol/administration and dosage”[Majr] AND (child OR children OR pediatr* OR paediatr*)

76 selected items

Records identified through database searching *n* = 76

Studies Included in this Review

Citation	Study Type
No studies answered the question	

Studies Not Included in this Review with Exclusion Rationale

Citation	Reason for exclusion
Abaya et al. (2019)	Continuous albuterol dosing
Battistini (2000)	Non-English
D’Vaz et al., (2019)	Dose not based on weight or age
Muchão et al. (2016)	High versus low dose
Parlar-Chun and Arnold (2021)	Continuous albuterol dosing
Polat, Saz, and Nursoy (2011)	Study on high dose Salbutamol
Ratnayake et al. (2016)	Dose not based on weight or age
Schuh et al. (1999)	Continuous albuterol dosing
Schuh et al. (2012)	Continuous albuterol dosing

Methods Used for Appraisal and Synthesis

^aRayyan is a web-based software used for the initial screening of titles and/or abstracts for this analysis (Ouzzani, Hammady, Fedorowicz & Elmagarmid, 2017).

^bThe Appraisal of Guidelines Research and Evaluation II (AGREE II) is an international instrument used to assess the quality and reporting of clinical practice guidelines for this analysis (Brouwers et al. 2010).

^cThe Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram depicts the process in which literature is searched, screened, and eligibility criteria is applied (Moher, Liberati, Tetzlaff, & Altman, 2009).

^aOuzzani, M., Hammady, H., Fedorowicz, Z., & Elmagarmid, A. (2016). Rayyan-a web and mobile app for systematic reviews. *Systematic Reviews*, 5(1), 210. Doi:10.1186/s13643-016-0384-4

^bBrouwers, M.C. et al. for the AGREE Next Steps Consortium. (2010) AGREE II: Advancing guideline development, reporting and evaluation in healthcare. *Canadian Medical Association Journal*, 182, E839-842. Retrieved from <https://www.agreetrust.org/wp-content/uploads/2017/12/AGREE-II-Users-Manual-and-23-item-Instrument-2009-Update-2017.pdf>

^cMoher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PloS Med* 6(7): e1000097. Doi:10.1371/journal.pmed1000097 **For more information, visit www.prisma-statement.org.**

Question Originator

H. Murphy, BHS, RRT AE-C

Medical Librarian Responsible for the Search Strategy

K. Swaggart, MLIS, AHIP

EBP Team or EBP Scholar’s Responsible for Analyzing the Literature

T. Bontrager, MSN, RN, CPEN

J. Dusin, MS, RD, LD, CPHQ
 J. Edwards, RN, MSN, CPEN
 K. Hess, PharmD
 H. Murphy, BHS RRT AE-C
 A. Wilson, BSN, RN, CPN


EBP Team Member Responsible for Reviewing, Synthesizing, and Developing this Document

J. Dusin, MS, RD, LD, CPHQ

Acronyms Used in this Document

Acronym	Explanation
AGREE II	Appraisal of Guidelines Research and Evaluation II
CAT	Critically Appraised Topic
EBP	Evidence Based Practice
EPR	Expert Panel Report
GINA	Global Initiative for Asthma
MDI	Metered dose inhaler
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analyses

Figure 1
Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)^c





Appendix D: Power Plan for Ambulatory Clinics

Being finalized, will add once available.

** These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.*



Appendix E: Quick Notes for Urgent Care

Being finalized, will add once available.

** These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.*

Appendix F: Power Plan for Emergency Department

+ Add to Phase Start: ... Duration: ...

\$	▼	Component	Status	Dose ...	Details
EDP Asthma CPG (Initiated Pending)					
Vital Signs/Monitoring					
<input type="checkbox"/>	<input checked="" type="checkbox"/>	Vital signs			
<input type="checkbox"/>	<input checked="" type="checkbox"/>	Blood Pressure			Upper Systolic Limit: 120, Lower Systolic Limit: 70, Upper Diastolic Limit: 80, Lower Diastolic Limit: 30, Upper MAP Limit: 90, Lower MAP Limit: 45
<input type="checkbox"/>	<input checked="" type="checkbox"/>	Temperature			
<input type="checkbox"/>	<input checked="" type="checkbox"/>	Oxygen/Pulse oximetry			Target Sat: >/= 90% (Standard), Lower alarm limit: 88, Upper alarm limit: 101
<input type="checkbox"/>	<input checked="" type="checkbox"/>	Cardiorespiratory monitor			Frequency: Continuous, RN to change limits Yes, Upper HR limit 150, Lower HR limit 60, Upper RR limit 50, Lower RR limit 12, Cardiorespiratory Leads 3
<input type="checkbox"/>	<input checked="" type="checkbox"/>	Height/Length			
Nutrition/Diet					
<input type="checkbox"/>	<input checked="" type="checkbox"/>	NPO diet			
Respiratory					
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Asthma Education by RT			
<input type="checkbox"/>	<input checked="" type="checkbox"/>	Asthma Action Plan			
<input type="checkbox"/>	<input checked="" type="checkbox"/>	Positive Expiratory Pressure (PEP)			
Consults/Therapy					
<input type="checkbox"/>	<input checked="" type="checkbox"/>	Consult to Child Life			T;N, Urgent
Laboratory					
<input type="checkbox"/>	<input checked="" type="checkbox"/>	COVID-19 Rapid RT PCR			
Miscellaneous					
<input type="checkbox"/>	<input checked="" type="checkbox"/>	Mild Asthma - PRAM Score 2 to 4			
<input type="checkbox"/>	<input checked="" type="checkbox"/>	Moderate Asthma - PRAM Score 5 to 7			
<input type="checkbox"/>	<input checked="" type="checkbox"/>	Severe Asthma - PRAM greater than 7			
<input type="checkbox"/>	<input checked="" type="checkbox"/>	Non Invasive Ventilation			

Mild Asthma Exacerbation Subphase:
Being finalized, will add once available.

Moderate Asthma Exacerbation Subphase:
Being finalized, will add once available.

Severe Asthma Exacerbation subphase:
Being finalized, will add once available.

* These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.

Appendix G: Power Plan for Pediatric Intensive Care

Component	Status	Dose ...	Details
PICU Status Asthmaticus (Initiated Pending)			
Admit/Transfer			
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Admit or Refer to Observation Status: Admit patient to inpatient services, Intensive Care, Pediatrics, Critical
Vital Signs/Monitoring			
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Vital signs q 2 hours
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Height/Length 1 time only On admission
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Weight 1 time only, on admission
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Weight Mon and Thurs
Nutrition/Diet			
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	NPO diet T;N, except oral medications
Nursing			
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Activity Bed rest-strict (no bathroom privileges)
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Intake and Output Strict
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Heparin flush for central and midlines (per CMH guidelines)
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Call Provider For increased respiratory distress or hypoxemia
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Call Provider For altered mental status
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Call Provider For temperature ≥ 38.5 C
<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Sequential compression device placement/assessment

Respiratory Orders with filtered order sentences

Respiratory			
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Oxygen/Pulse oximetry Target Sat: $\geq 90\%$ (Standard), Frequency: Continuous, Lower alarm limit: 88, Upper alarm limit: 101, Indications: Critically ill
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Respiratory Care Plan
<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	albuterol (albuterol continuous for *NON*mechanically ventilated patients) 15 milligrams per hour, NEB, per protocol, PRN Wheezing or Cough, For p... Albuterol 0.5% = 5mg/ml
<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	albuterol (albuterol continuous for invasive and non-invasive mechanically ventilated patients) 15 milligrams per hour, NEB, per protocol, PRN Wheezing or Cough, For p... **mechanically ventilated: use 60 ml bag ** Albuterol 0.5% = 5mg/ml
<input type="checkbox"/> Filtered Order Sentences			
15 milligrams per hour, NEB, per protocol, PRN Wheezing or Cough, For patients 20 kg or greater Greater Than or Equal To 20 kg			
20 milligrams per hour, NEB, per protocol, PRN Wheezing or Cough			
10 milligrams per hour, NEB, per protocol, PRN Wheezing or Cough, For patients < 20 kg Less Than 20 kg			
<input type="checkbox"/> Filtered Order Sentences			
15 milligrams per hour, NEB, per protocol, PRN Wheezing or Cough, For patients 20 kg or greater **mechanically ventilated: use 60 ml bag ** Greater Than or Equal To 20 kg			
20 milligrams per hour, NEB, per protocol, PRN Wheezing or Cough			
10 milligrams per hour, NEB, per protocol, PRN Wheezing or Cough, For patients < 20 kg **mechanically ventilated: use 60 ml bag ** Less Than 20 kg			

* These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.

Consults/Therapy and Labs

Consults/Therapy		
<input checked="" type="checkbox"/>		High Risk Asthma is > or = 4 ED/UCC/inpatient episodes in the past 12 months (including current visit) OR current PICU admission. (admission resulting from ED/UCC visit counts as one episode)
<input checked="" type="checkbox"/>		High Risk Asthma Initiated Pen...
<input checked="" type="checkbox"/>		For Asthma Education only, please page the respiratory therapist or Asthma Coordinator (458-3546). Asthma Class
Laboratory		
<input type="checkbox"/>		Admission Lab
<input type="checkbox"/>		CBC w/Differential Blood, Stat collect
<input type="checkbox"/>		Basic Metabolic Panel Blood, Stat collect
<input type="checkbox"/>		hCG Qual Urine Urine, Stat collect, T;N, Time Change Allowed Yes
<input type="checkbox"/>		Routine Lab
<input type="checkbox"/>		Basic Metabolic Panel Blood, Timed Study/Recurring collect, Time Change Allowed Yes, q24h Std (0400). Nurse collect
<input type="checkbox"/>		If on aminophylline drip Theophylline (Aminophylline) Level Blood, Timed Study/Recurring collect, T;N, Time Change Allowed Yes, Nurse collect

Continuous Medications/Fluids

Continuous Medications/Fluids		
<input type="checkbox"/>		sodium chloride 0.9% (normal saline) Order sentences below are for Oncology Powerchart only
<input type="checkbox"/>		DSW with 0.45% NaCl and KCl 20 mEq/l 1,000
<input type="checkbox"/>		Aminophylline load and drip aminophylline T;N, 6 mg/kg, IV infuse via pump over 30 min
<input type="checkbox"/>		Aminophylline in DSW 10 mg/mL, continous (standard)
<input type="checkbox"/>		Order IV Placement and saline flush for peripheral lines.
<input checked="" type="checkbox"/>		IV placement
<input type="checkbox"/>		Heparin flush for central and midlines (per CMH guidelines)
<input type="checkbox"/>		IV + PO
<input checked="" type="checkbox"/>		Discontinue IVF from previous encounter

Medications

Medications		
<input checked="" type="checkbox"/>		acetaminophen 10 mg/kg, PO, q4hr, PRN Fever
<input type="checkbox"/>		acetaminophen 10 mg/kg, Per Rectum, Supp, q4hr, PRN Fever
<input type="checkbox"/>		albuterol (albuterol HFA 90 mcg/inh inhalation aerosol) 4 puff, Inhaled, per protocol, PRN Wheezing or Cough May use 2- 8 puffs per dose per respiratory care plan. AT HOME: Use as d...
<input type="checkbox"/>		albuterol (albuterol 2.5 mg/3 mL (0.083%) inhalation solution) 3 mL, NEB, per protocol, PRN Wheezing
<input type="checkbox"/>		ipratropium (ipratropium 17 mcg/inh inhaler) 1 puff, Inhaled, q8hr, Wheezing
<input type="checkbox"/>		ipratropium (ipratropium 500 mcg/2.5 mL inhalation solution) 1,500 mcg, NEB, q8hr Give with continuous albuterol
<input type="checkbox"/>		magnesium sulfate 50 mg/kg, IV, 1 time only
<input type="checkbox"/>		fluticasone (Flovent HFA 44 mcg/inh inhalation aerosol with adapter) 1 puff, Inhaled, BID
<input type="checkbox"/>		fluticasone (Flovent HFA 110 mcg/inh inhalation aerosol with adapter) 1 puff, Inhaled, BID
<input type="checkbox"/>		fluticasone (Flovent HFA 220 mcg/inh inhalation aerosol with adapter) 1 puff, Inhaled, BID
<input type="checkbox"/>		prednisolONE (prednisolONE (as sodium phosphate) 15 mg/5 mL oral liquid) 30 mg, PO, BID
<input type="checkbox"/>		predniSONE 30 mg, PO, BID
<input checked="" type="checkbox"/>		methylPREDNISolone 0.5 mg/kg, IV, q6hr
<input type="checkbox"/>		famotidine (famotidine injectable) 20 mg, IV, q12hr
<input type="checkbox"/>		pantoprazole (pantoprazole IV) 1 mg/kg, IV, q24hr
<input type="checkbox"/>		Asthma Action Plan

* These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.

Appendix H: Power Plan for Hospital Medicine

Component	Status	Dose ...	Details
Inpatient Asthma CPG (Planned Pending)			
Admit/Transfer	<input checked="" type="checkbox"/>		This Powerplan is intended for otherwise healthy patients >60 days of age with suspected asthma exacerbation.
	<input checked="" type="checkbox"/>		Admit or Refer to Observation
Seasonal Immunizations	<input type="checkbox"/>		influenza virus vaccine, inactivated 0.5 mL, IM, Unscheduled, 1 dose(s) obtain consent prior to administration.
Vital Signs/Monitoring			
	<input checked="" type="checkbox"/>		Vital signs Select an order sentence
	<input checked="" type="checkbox"/>		Weight 1 time only On admission
	<input checked="" type="checkbox"/>		Height/Length 1 time only On admission
Nutrition/Diet			
	<input type="checkbox"/>		Regular diet for age
	<input type="checkbox"/>		NPO Diet Instructions
	<input type="checkbox"/>		Diets
Nursing			
	<input checked="" type="checkbox"/>		Intake and Output Measured
	<input checked="" type="checkbox"/>		Activity As tolerated/ad lib
	<input type="checkbox"/>		Consider isolation order for patients that have URI symptoms
	<input type="checkbox"/>		Isolation Contact / Droplet Patient should be in isolation if URI symptoms are present.
	<input checked="" type="checkbox"/>		PEWS Baseline Assessment
	<input type="checkbox"/>		IV placement
	<input type="checkbox"/>		IV + PO
	<input type="checkbox"/>		Saline lock (Saline lock IV line when taking adequate PO)
	<input type="checkbox"/>		Sequential compression device placement/assessment (SCD Placement/assessment)
Respiratory			
	<input checked="" type="checkbox"/>		Asthma Action Plan
	<input checked="" type="checkbox"/>		Respiratory Care Plan Beta-Agonist Indications: History of asthma Use PEP at discretion of respiratory therapist
	<input checked="" type="checkbox"/>		Oxygen/Pulse oximetry Frequency: Intermittent q4, Target Sat: >= 90% (Standard), Lower alarm limit: 88, Upper alarm limit: 101
Consults/Therapy			
	<input checked="" type="checkbox"/>		High Risk Asthma is > or = 4 ED/UCC/inpatient episodes in the past 12 months (including current visit), current PICU admission, history of intubation AND/OR cardiac arrest due to asthma. (admission resulting from ED/UCC visit counts as one episode)
	<input type="checkbox"/>		High Risk Asthma
	<input type="checkbox"/>		For Asthma Education only, please page the respiratory therapist or Asthma Coordinator (458-3546).
	<input checked="" type="checkbox"/>		Asthma Class
	<input type="checkbox"/>		Consider Allergy/Immunology Consult for patients with food allergies or severe eczema
	<input type="checkbox"/>		Consult to Allergy/Immunology
	<input type="checkbox"/>		Consult to Pulmonology
	<input type="checkbox"/>		Consult to Social Work
	<input type="checkbox"/>		Environmental Health Home Education Referral
Continuous Medications/Fluids			
	<input type="checkbox"/>		sodium chloride 0.9% (normal saline)
	<input type="checkbox"/>		Discontinue IVF from previous encounter
Medications			
	<input type="checkbox"/>		See CPG guidelines for Mild, Moderate, Severe considerations
	<input type="checkbox"/>		Mild Asthma Exacerbation
	<input type="checkbox"/>		Moderate Asthma Exacerbation
	<input type="checkbox"/>		Severe Asthma Exacerbation
	<input type="checkbox"/>		Severe Asthma Exacerbation
Controllers			
	<input type="checkbox"/>		fluticasone (Flovent HFA 44 mcg/inh inhalation aerosol with adapter) 2 puff, Inhaled, BID
	<input type="checkbox"/>		fluticasone (Flovent HFA 110 mcg/inh inhalation aerosol with adapter) 2 puff, Inhaled, BID
	<input type="checkbox"/>		fluticasone (Flovent HFA 220 mcg/inh inhalation aerosol with adapter) 2 puff, Inhaled, BID
	<input type="checkbox"/>		budesonide (budesonide 0.5 mg/2 mL inhalation suspension) 0.5 mg, NEB, BID
	<input type="checkbox"/>		budesonide (budesonide 1 mg/2 mL inhalation suspension) 1 mg, NEB, qDay
	<input type="checkbox"/>		budesonide (budesonide 0.25 mg/2 mL inhalation suspension) 0.25 mg, NEB, BID
	<input type="checkbox"/>		fluticasone/salmeterol (fluticasone-salmeterol 45 mcg-21 mcg inhalation aerosol with adapter) 2 puff, Inhaled, BID
	<input type="checkbox"/>		fluticasone/salmeterol (fluticasone-salmeterol 115 mcg-21 mcg inhalation aerosol with adapter) 2 puff, Inhaled, BID
	<input type="checkbox"/>		fluticasone/salmeterol (fluticasone-salmeterol 230 mcg-21 mcg inhalation aerosol with adapter) 2 puff, Inhaled, BID
	<input type="checkbox"/>		montelukast (montelukast 4 mg oral granule) 4 mg, PO, qDay
	<input type="checkbox"/>		montelukast (montelukast 4 mg oral tablet, chewable) 4 mg, PO, qDay
	<input type="checkbox"/>		montelukast (montelukast 5 mg oral tablet, chewable) 5 mg, PO, qDay
	<input type="checkbox"/>		montelukast (montelukast 10 mg oral tablet) 10 mg, PO, qDay

* These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.

Mild Asthma Exacerbation Subphase:

Component	Status	Dose ...	Details
Inpatient Asthma CPG, Mild Asthma Exacerbation (Planned Pending)			
Medications			
albuterol (albuterol HFA 90 mcg/inh inhalation aerosol)		8 puff, Inhaled, per protocol AT HOME: Use as directed per discharge	
albuterol (albuterol 2.5 mg/3 mL (0.083%) inhalation solution)		4 puff, Inhaled, per protocol	Greater Than or Equal To 20 kg Less Than 20 kg
Steroids			
Consider steroids if >= 2 albuterol doses are required.			
dexAMETHasone		12 mg, PO, 1 time only Max Dose: 12mg/dose	
prednisolONE (prednisolONE (as sodium phosphate) 15 mg/5 mL oral liquid)		60 mg, PO, qDay	
predniSONE		60 mg, PO, qDay MAX DOSE: 60 mg	

Dexamethasone order sentences:

Steroids			
Consider steroids if >= 2 albuterol doses are required.			
dexAMETHasone		12 mg, PO, 1 time only Max Dose: 12mg/dose	
prednisolONE (prednisolONE (as sodium phosphate) 15 mg/5 mL oral liquid)		60 mg, PO, qDay	
predniSONE		12 mg, PO, 1 time only 0.6 mg/kg, PO, 1 time only	Greater Than or Equal To 20 kg Less Than 20 kg

Prednisolone order sentences:

prednisolONE (prednisolONE (as sodium phosphate) 15 mg/5 mL oral liquid)		60 mg, PO, qDay	
predniSONE		60 mg, PO, qDay	Greater Than or Equal To 30 kg Less Than 30 kg

Prednisone order sentences:

predniSONE		60 mg, PO, qDay MAX DOSE: 60 mg	
		60 mg, PO, qDay	Greater Than or Equal To 20 kg Less Than 30 kg

Moderate Asthma Exacerbation Subphase:

Component	Status	Dose ...	Details
Inpatient Asthma CPG, Moderate Asthma Exacerbation (Planned Pending)			
Medications			
albuterol (albuterol HFA 90 mcg/inh inhalation aerosol)		8 puff, Inhaled, per protocol AT HOME: Use as directed per discharge instructions.	
Combined NEB of Albuterol and Ipratropium : 1 hour of continuous albuterol with or without 1500 mcg of ipratropium for the initial			
albuterol (albuterol continuous for "NON"mechanically ventilated patients)		8 puff, Inhaled, per protocol	Greater Than or Equal To 20 kg Less Than 20 kg
ipratropium (ipratropium 0.02% inhalation solution)		4 puff, Inhaled, per protocol	
Steroids			
dexAMETHasone		12 mg, PO, 1 time only Max Dose: 12mg/dose	
prednisolONE (prednisolONE (as sodium phosphate) 15 mg/5 mL oral liquid)		Select an order sentence	
predniSONE		60 mg, PO, qDay MAX DOSE: 60 mg	
Other Medications			
magnesium sulfate (magnesium sulfate bolus 60 mg/mL (peripheral line))		2,000 mg, IV, 1 time only Max Dose: 2 grams. Run in over 20 minutes. Check blood pressure q5min once infusion has started.	
Analgesics			
lidocaine/sodium bicarbonate (I-Tip with buffered lidocaine 0.9%)		0.2 mL, Intradermal, Unscheduled, PRN Needle Sticks, 1 dose(s)	

Combined neb order sentences:

Combined NEB of Albuterol and Ipratropium : 1 hour of continuous albuterol with or without 1500 mcg of ipratropium for the initial care of the patient in moderate patient.			
albuterol (albuterol continuous for "NON"mechanically ventilated patients)		15 milligrams per hour, NEB, per protocol, PRN Wheezing or Cough Albuterol 0.5% = 5mg/ml	
ipratropium (ipratropium 0.02% inhalation solution)		10 milligrams per hour, NEB, per protocol, PRN Wheezing or Cough	Greater Than or Equal To 20 kg Less Than 20 kg

Magnesium sulfate order sentences:

magnesium sulfate (magnesium sulfate bolus 60 mg/mL (peripheral line))		2,000 mg, IV, 1 time only Max Dose: 2 grams. Run in over 20 minutes. Check blood pressure q5min once infusion has started.	
Analgesics			
lidocaine/sodium bicarbonate (I-Tip with buffered lidocaine 0.9%)		2,000 mg, IV, 1 time only	Greater Than or Equal To 40 kg Less Than 40 kg

Prednisone order sentences:

predniSONE		60 mg, PO, qDay MAX DOSE: 60 mg	
Other Medications			
magnesium sulfate (magnesium sulfate bolus 60 mg/mL (peripheral line))		60 mg, PO, qDay	Greater Than or Equal To 30 kg Less Than 30 kg
Analgesics			
lidocaine/sodium bicarbonate (I-Tip with buffered lidocaine 0.9%)		2 mg/kg, PO, qDay	Less Than 30 kg

* These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.

Severe Asthma Exacerbation subphase:

Component	Status	Dose ...	Details
Inpatient Asthma CPG, Severe Asthma Exacerbation (Planned Pending)			
Medications			
<input type="checkbox"/>		albuterol (albuterol continuous for *NON*mechanically ventilated patients)	15 milligrams per hour, NEB, per protocol, PRN Wheezing or Cough Albuterol 0.5% = 5mg/ml
<input type="checkbox"/>		albuterol (albuterol continuous for invasive and non-invasive mechanically ventilated patients)	<input type="checkbox"/> Filtered Order Sentences
<input type="checkbox"/>		ipratropium (ipratropium 0.02% inhalation solution)	15 milligrams per hour, NEB, per protocol, PRN Wheezing or Cough Greater Than or Equal To 20 kg 10 milligrams per hour, NEB, per protocol, PRN Wheezing or Cough Less Than 20 kg
<input type="checkbox"/>		methyIPREDNISolone	MAX DOSE: 60mg
<input type="checkbox"/>		magnesium sulfate (magnesium sulfate bolus 60 mg/mL (peripheral line))	2,000 mg, IV, 1 time only Max Dose: 2 grams. Run in over 20 minutes. Check blood pressure q5min once infusion has started.
<input type="checkbox"/>		EPINEPHrine (EPINEPHrine 1 mg/mL injection)	0.5 mg, IM, 1 time only [Epinephrine 1 mg/mL] MAX DOSE: 0.5mg
<input type="checkbox"/>		EPINEPHrine (EpiPen JR Auto-Injector)	Select an order sentence
<input type="checkbox"/>		EPINEPHrine (EpiPen Auto-Injector)	0.3 mg, IM, 1 time only, PRN Anaphylaxis, EpiPen (For patients greater than or equal to 25 kg) For patients greater than or equal to 25 kg
Analgesics			
<input type="checkbox"/>		lidocaine/sodium bicarbonate (J-Tip with buffered lidocaine 0.9%)	0.2 mL, Intradermal, Unscheduled, PRN Needle Sticks, 1 dose(s)
<input type="button" value="Return to Inpatient Asthma CPG"/>			

** These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.*

Appendix I: AGREE II Assessment for Children's Mercy Hospital's Asthma CPG

AGREE II^a Summary for this Clinical Practice Guideline*

Domain	Percent Agreement
Scope and purpose	100%
Stakeholder involvement	92%
Rigor of development	99%
Clarity and presentation	100%
Applicability	98%
Editorial independence	100%
Reviewer's recommendation for guideline use	Adopt the utilization of this guideline

*Note: This assessment reflects the views obtained from one external clinician and one internal clinician.

** These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different, and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly, these guidelines should guide care with the understanding that departures from them may be required at times.*